



2026/2024

11.9.2026

COMMISSION IMPLEMENTING REGULATION (EU) 2026/2024

of 10 September 2026

amending Implementing Regulation (EU) 2024/2750 as regards administrative and minor changes to the Union authorisation of the biocidal product family 'LANXESS CMIT/MIT biocidal product family'

(Text with EEA relevance)

THE EUROPEAN COMMISSION,

Having regard to the Treaty on the Functioning of the European Union,

Having regard to Regulation (EU) No 528/2012 of the European Parliament and of the Council of 22 May 2012 concerning the making available on the market and use of biocidal products ⁽¹⁾, and in particular Article 50(2) thereof,

Whereas:

- (1) On 25 October 2024, Commission Implementing Regulation (EU) 2024/2750 ⁽²⁾ granted a Union authorisation, under number EU-0031652-0000, to Lanxess Deutschland GmbH for the making available on the market and use of the biocidal product family 'LANXESS CMIT/MIT biocidal product family'. The Annex to that Implementing Regulation provides the summary of product characteristics ('SPC') for that biocidal product family.
- (2) Commission Implementing Regulation (EU) No 354/2013 ⁽³⁾ lays down the procedural rules for the different categories of changes referred to in Article 50(3) of Regulation (EU) No 528/2012.
- (3) On 24 July 2025, Lanxess Deutschland GmbH submitted to the European Chemicals Agency ('the Agency'), in accordance with Article 12(1) of Implementing Regulation (EU) No 354/2013, an application for minor changes to the Union authorisation for the biocidal product family 'LANXESS CMIT/MIT biocidal product family', as referred to in Title 2 of the Annex to that Regulation. The application was recorded in the Register for Biocidal Products under case number BC-HT109132-36. The proposed changes to that authorisation concern the extension of the shelf life of products in meta-SPC 1 to 6.
- (4) On 27 October 2025 and 6 March 2026, Lanxess Deutschland GmbH submitted to the Agency, in accordance with Article 11(1) of Implementing Regulation (EU) No 354/2013, notifications of administrative changes to the Union authorisation for the biocidal product family 'LANXESS CMIT/MIT biocidal product family', as referred to in Title 1 of the Annex to that Regulation. The notifications were recorded in the Register for Biocidal Products under case numbers BC-NY111447-09 and BC-NG116317-45. The notified proposed changes to that authorisation concern the addition of product manufacturers and trade names.
- (5) On 6 November 2025, the Agency submitted, in accordance with Article 12(4) of Implementing Regulation (EU) No 354/2013, an opinion ⁽⁴⁾ on the applied minor changes to the Union authorisation for the biocidal product family 'LANXESS CMIT/MIT biocidal product family' together with a revised SPC and a revised assessment report. The opinion concludes that the proposed changes are minor changes as defined in Article 3(1), point (ab), of Regulation (EU) No 528/2012 and as specified in Title 2 of the Annex to Implementing Regulation (EU) No 354/2013, and that after the implementation of the changes, the conditions of Article 19 of Regulation (EU) No 528/2012 will still be met.

⁽¹⁾ OJ L 167, 27.6.2012, p. 1, ELI: <http://data.europa.eu/eli/reg/2012/528/oj>.

⁽²⁾ Commission Implementing Regulation (EU) 2024/2750 of 25 October 2024 granting a Union authorisation for the biocidal product family LANXESS CMIT/MIT biocidal product family in accordance with Regulation (EU) No 528/2012 of the European Parliament and of the Council (OJ L, 2024/2750, 28.10.2024, ELI: http://data.europa.eu/eli/reg_impl/2024/2750/oj).

⁽³⁾ Commission Implementing Regulation (EU) No 354/2013 of 18 April 2013 on changes of biocidal products authorised in accordance with Regulation (EU) No 528/2012 of the European Parliament and of the Council (OJ L 109, 19.4.2013, p. 4, ELI: http://data.europa.eu/eli/reg_impl/2013/354/oj).

⁽⁴⁾ Biocidal Products Committee (BPC) opinion on the minor change to the Union authorisation of the biocidal product family 'LANXESS CMIT/MIT biocidal product family' of 6 November 2025, ECHA/BPC/505/2025 <https://echa.europa.eu/opinions-on-union-authorisation>.

- (6) On 26 November 2025 and 31 March 2026, the Agency submitted to the Commission, in accordance with Article 11(3) of Implementing Regulation (EU) No 354/2013, opinions ⁽⁹⁾ on the notified administrative changes to the Union authorisation for the biocidal product family 'LANXESS CMIT/MIT biocidal product family', together with a revised SPC. In these opinions, the Agency concludes that the proposed changes are administrative changes as defined in Article 3(1), point (aa), of Regulation (EU) No 528/2012 and as specified in Title 1, Sections 1 and 2, of the Annex to Implementing Regulation (EU) No 354/2013, and that, after the implementation of the changes, the conditions of Article 19 of Regulation (EU) No 528/2012 will still be met.
- (7) On 31 March 2026, the Agency transmitted to the Commission the revised summary of the product characteristics of the Union authorisation for the biocidal product family 'LANXESS CMIT/MIT biocidal product family' in all official languages of the Union, covering all the administrative changes and minor changes notified and applied for, in accordance with Article 11(6) and Article 12(6) of Implementing Regulation (EU) No 354/2013.
- (8) The Commission concurs with the opinions of the Agency and therefore considers it appropriate to amend the Union authorisation for the biocidal product family 'LANXESS CMIT/MIT biocidal product family' to introduce the administrative and minor changes requested by Lanxess Deutschland GmbH.
- (9) In order to enhance clarity and to ease the access of users and interested parties to the consolidated version of the summary of product characteristics which is to be published by the Agency, the Annex to Implementing Regulation (EU) 2024/2750 should be replaced in its entirety.
- (10) Implementing Regulation (EU) 2024/2750 should therefore be amended accordingly,

HAS ADOPTED THIS REGULATION:

Article 1

The Annex to Implementing Regulation (EU) 2024/2750 is replaced by the text set out in the Annex to this Regulation.

Article 2

This Regulation shall enter into force on the twentieth day following that of its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels, 10 September 2026.

For the Commission
The President
Ursula VON DER LEYEN

⁽⁹⁾ ECHA Opinion No UAD-C-1874461-24-00/F of 26 November 2025 and Opinion No UAD-C-1912183-29-00/F of 31 March 2026 on administrative changes to the Union authorisation of the biocidal product family 'LANXESS CMIT/MIT biocidal product family' <https://echa.europa.eu/opinions-on-union-authorisation>.

ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT FAMILY

LANXESS CMIT/MIT biocidal product family

Product type(s)

PT06: Preservatives for products during storage

PT11: Preservatives for liquid-cooling and processing systems

PT12: Slimicides

PT13: Working or cutting fluid preservatives

Authorisation number EU-0031652-0000

R4BP asset number EU-0031652-0000

PART I

FIRST INFORMATION LEVEL

1. ADMINISTRATIVE INFORMATION

1.1. Family name

Name	LANXESS CMIT/MIT biocidal product family
------	--

1.2. Product type(s)

Product type(s)	PT06: Preservatives for products during storage PT11: Preservatives for liquid-cooling and processing systems PT12: Slimicides PT13: Working or cutting fluid preservatives
-----------------	--

1.3. Authorisation holder

Name and address of the authorisation holder	Name	Lanxess Deutschland GmbH
	Address	Kennedyplatz 1 50569 Köln Germany
Authorisation number	EU-0031652-0000	
R4BP asset number	EU-0031652-0000	
Date of the authorisation	17.11.2024	
Expiry date of the authorisation	31.10.2034	

1.4. **Manufacturer(s) of the product**

Name of manufacturer	Lanxess Deutschland GmbH, BU Material Protection Products
Address of manufacturer	Kennedyplatz 1 D-50569 Köln Germany
Location of manufacturing sites	Lanxess Deutschland GmbH, BU Material Protection Products site 1 Rheinuferstraße 7-9 47829 Krefeld Germany
Name of manufacturer	Lanxess Pte. Ltd.
Address of manufacturer	16, Joo Koon Crescent 629018 Singapore Singapore
Location of manufacturing sites	Lanxess Pte. Ltd. site 1 16, Joo Koon Crescent 629018 Singapore Singapore
Name of manufacturer	LANXESS Chemical (China) Co., Ltd
Address of manufacturer	No. 318, Huanghai Road 213127 Xinbei District, Changzhou, Jiangsu Province China
Location of manufacturing sites	LANXESS Chemical (China) Co., Ltd site 1 No. 318, Huanghai Road 213127 Xinbei District, Changzhou, Jiangsu Province China
Name of manufacturer	LANXESS Corporation
Address of manufacturer	Neville Island, 3499 Grand Avenue 15225 Pittsburgh, PA United States (the)
Location of manufacturing sites	LANXESS Corporation site 1 Neville Island, 3499 Grand Avenue 15225 Pittsburgh, PA United States (the)
Name of manufacturer	LANXESS India Pvt. Ltd.
Address of manufacturer	Jhagadia Industrial Estate, Plot No 748/2/A, GIDC 393110 District Bharuch, Jhagadia India
Location of manufacturing sites	LANXESS India Pvt. Ltd. site 1 Plot No 748/2/A, GIDC 393110 District Bharuch, Jhagadia India
Name of manufacturer	Vera Chimie Productions
Address of manufacturer	Zone Industrielle du Broteau, 2 Rue Du Broteau 69540 Irigny France
Location of manufacturing sites	Vera Chimie Productions site 1 Zone Industrielle du Broteau, 2 Rue Du Broteau 69540 Irigny France

Name of manufacturer	Kemi-tech ApS
Address of manufacturer	Mosetoften 14 8722 Hedensted Denmark
Location of manufacturing sites	Kemi-tech ApS site 1 Mosetoften 14 8722 Hedensted Denmark
Name of manufacturer	Servophil AG
Address of manufacturer	Bösch 73 6331 Hünenberg Switzerland
Location of manufacturing sites	Servophil AG site 1 Bösch 73 6331 Hünenberg Switzerland
Name of manufacturer	Theseo Deutschland GmbH
Address of manufacturer	Kolpingstrasse 4 49835 Wietmarschen Germany
Location of manufacturing sites	Theseo Deutschland GmbH Kolpingstrasse 4 49835 Wietmarschen Germany
Name of manufacturer	URSA-Chemie GmbH
Address of manufacturer	Am Alten Galgen 14 56410 Montabaur Germany
Location of manufacturing sites	URSA-Chemie GmbH Am Alten Galgen 14 56410 Montabaur Germany
Name of manufacturer	Cellwater Lda
Address of manufacturer	Rua dos Armazéns Nº 7 2330-210 Entroncamento Portugal
Location of manufacturing sites	Cellwater Lda Rua dos Armazéns Nº 7 2330-210 Entroncamento Portugal
Name of manufacturer	TÉCNICAS DE DOSIFICACIÓN Y FILTRACIÓN, S.L.-TDF
Address of manufacturer	C/ Roma, 13 28970 Humanes de Madrid Spain
Location of manufacturing sites	TÉCNICAS DE DOSIFICACIÓN Y FILTRACIÓN, S.L.-TDF C/ Roma, 13 28970 Humanes de Madrid Spain
Name of manufacturer	ANDALUZA DE TRATAMIENTOS INDUSTRIALES, S.L.-ANTRAIN
Address of manufacturer	Camino Huerta Santa Cruz, Nº29 - Pol. Ind, Huerta Santa Cruz 29004 Malaga Spain
Location of manufacturing sites	ANDALUZA DE TRATAMIENTOS INDUSTRIALES, S.L.-ANTRAIN Camino Huerta Santa Cruz, Nº29 - Pol. Ind, Huerta Santa Cruz 29004 Malaga Spain

Name of manufacturer	OTARI Ingeniería del Agua, S.L.
Address of manufacturer	Pol. Ind. Arretxe-Ugalde - Ezurriki Kalea, 31 20305 Irun Spain
Location of manufacturing sites	OTARI Ingeniería del Agua, S.L. Pol. Ind. Arretxe-Ugalde - Ezurriki Kalea, 31 20305 Irun Spain

Name of manufacturer	QUIMICA PARA LAS FASES DE AGUA, S.L
Address of manufacturer	Pol. Alcoz Alto 16 50410 Cuarte de Huerva Spain
Location of manufacturing sites	QUIMICA PARA LAS FASES DE AGUA, S.L Pol. Alcoz Alto 16 50410 Cuarte de Huerva Spain

Name of manufacturer	ODYL, S.A.
Address of manufacturer	C/Maresme, 19 - Pol. Ind. Can Bernades-Subira 08130 Santa Perpetua De Mogoda Spain
Location of manufacturing sites	ODYL, S.A. C/Maresme, 19 - Pol. Ind. Can Bernades-Subira 08130 Santa Perpetua De Mogoda Spain

Name of manufacturer	Lamirsa Solutions, S.L.
Address of manufacturer	Pol. Ind. Can Parellada - C/ Apolo, 46 08228 Terrassa Spain
Location of manufacturing sites	Lamirsa Solutions, S.L. Pol. Ind. Can Parellada - C/ Apolo, 46 08228 Terrassa Spain

Name of manufacturer	Kurita Europe GmbH
Address of manufacturer	Theodor-Heuss-Anlage 2 68165 Mannheim Germany
Location of manufacturing sites	Kurita Europe GmbH Giulinistrasse 2 67065 Mannheim Germany

Name of manufacturer	Tintometer GmbH, Lovibond® Water Testing
Address of manufacturer	Schleefstrasse 8-12 44287 Dortmund Germany
Location of manufacturing sites	Tintometer GmbH, Lovibond® Water Testing Schleefstrasse 8-12 44287 Dortmund Germany

1.5. **Manufacturer(s) of the active substance(s)**

Active substance	C(M)IT/MIT (3:1)
Name of manufacturer	Dalian Bio-Chem Co., Ltd.
Address of manufacturer	10F, R&F Center, No. 6 Gangxing Road, Zhongshan District 116001 Dalian China
Location of manufacturing sites	Dalian Bio-Chem Co., Ltd. site 1 Dalian Songmudao Chemical Industry Zone, Puwan New District, 116308 Dalian, Liaoning China

2. **PRODUCT FAMILY COMPOSITION AND FORMULATION**2.1. **Qualitative and quantitative information on the composition of the family**

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		1 - 24,2 % (w/w)

2.2. **Type(s) of formulation**

Formulation type(s)	AL Any other liquid
---------------------	---------------------

PART II

SECOND INFORMATION LEVEL META SPC(S)

1. **META SPC 1 ADMINISTRATIVE INFORMATION**1.1. **Meta SPC 1 identifier**

Identifier	Meta SPC: Meta SPC 1
------------	----------------------

1.2. **Suffix to the authorisation number**

Number	1-1
--------	-----

1.3. **Product type(s)**

Product type(s)	PT06: Preservatives for products during storage PT11: Preservatives for liquid-cooling and processing systems PT12: Slimicides PT13: Working or cutting fluid preservatives
-----------------	---

2. META SPC 1 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 1

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		24,2 - 24,2 % (w/w)

2.2. Type(s) of formulation of the meta SPC 1

Formulation type(s)	AL Any other liquid
---------------------	---------------------

3. HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 1

Hazard statements	H290: May be corrosive to metals. H302: Harmful if swallowed. H311: Toxic in contact with skin. H314: Causes severe skin burns and eye damage. H317: May cause an allergic skin reaction. H332: Harmful if inhaled. H410: Very toxic to aquatic life with long lasting effects. EUH071: Corrosive to the respiratory tract.
Precautionary statements	P234: Keep only in original packaging. P260: Do not breathe vapours. P260: Do not breathe dust. P260: Do not breathe gas. P260: Do not breathe mist. P260: Do not breathe spray. P260: Do not breathe fume. P270: Do not eat, drink or smoke when using this product. P271: Use only outdoors or in a well-ventilated area. P272: Contaminated work clothing should not be allowed out of the workplace. P273: Avoid release to the environment. P280: Wear protective gloves. P280: Wear protective clothing. P280: Wear eye protection. P280: Wear face protection. P301+P330+P331: IF SWALLOWED: rinse mouth. Do NOT induce vomiting. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water [or shower]. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor.

	P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P304+P340: IF INHALED: Remove person to fresh air and keep comfortable for breathing. P312: Call a POISON CENTER/doctor if you feel unwell. P361+P364: Take off immediately all contaminated clothing and wash it before reuse. P405: Store locked up. P406: Store in a corrosion-resistant container with a resistant inner liner. P501: Dispose of contents in accordance with all local, regional, national and international regulations. P501: Dispose of container in accordance with all local, regional, national and international regulations. P403+P233: Store in a well-ventilated place. Keep container tightly closed. P391: Collect spillage. P390: Absorb spillage to prevent material damage.
--	---

4. AUTHORISED USE(S) OF THE META SPC

4.1. Use description

Table 1

In-can preservation of washing and cleaning fluids (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Washing and cleaning fluids
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product has to be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 26,4 - 107,1 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected in such a way that a pure CMIT/MIT concentration of 3,7 to less than 15 ppm is maintained. Number and timing of application: Frequency: Single application

Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.1.1. *Use-specific instructions for use*

See general directions for use.

4.1.2. *Use-specific risk mitigation measures*

See general directions for use.

4.1.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.1.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.1.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.2. **Use description**

Table 2

In-can preservation of paints and coatings (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Paints and coatings
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product has to be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.

Application rate(s) and frequency	Application Rate: 79,3 – 107,1 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 11,1 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.2.1. *Use-specific instructions for use*

See general directions for use.

4.2.2. *Use-specific risk mitigation measures*

See general directions for use.

4.2.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.2.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.2.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.3. **Use description**

Table 3

In-can preservation of fluids used in paper production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in paper production (e.g. pigment slurries)

Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 52,9 – 214,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.3.1. *Use-specific instructions for use*

See general directions for use.

4.3.2. *Use-specific risk mitigation measures*

See general directions for use.

4.3.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.3.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.3.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.4. **Use description**

Table 4

In-can preservation of fluids used in textile production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -

Field(s) of use	indoor use Fluids used in textile production
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 26,4 – 214,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.4.1. Use-specific instructions for use

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

4.4.2. Use-specific risk mitigation measures

See general directions for use.

4.4.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.4.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.4.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.5. Use description

Table 5

In-can preservation of fluids used in leather production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in leather production
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product has to be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 52,9 – 214,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.5.1. *Use-specific instructions for use*

See general directions for use.

4.5.2. *Use-specific risk mitigation measures*

See general directions for use.

4.5.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.5.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.5.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.6. **Use description**

Table 6

In-can preservation of glues and adhesives (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Glues and adhesives
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 26,4 – 214,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 30 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.6.1. Use-specific instructions for use

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

See general directions for use.

4.6.2. Use-specific risk mitigation measures

See general directions for use.

4.6.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.6.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.6.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.7. Use description

Table 7

In-can preservation of concrete additives and building material such as fillers/sealants, plasters and wax emulsions (PT 6) for indoor use

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Concrete additives and building material such as fillers/sealants, plasters and wax emulsions intended for indoor use
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 52,8 - 107,1 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into product matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.7.1. *Use-specific instructions for use*

See general directions for use.

4.7.2. *Use-specific risk mitigation measures*

See general directions for use.

4.7.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.7.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.7.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.8. Use description

Table 8

In-can preservation of polymer dispersions/emulsions intended for use in paints and coatings, glues and adhesives, building materials and fluids used in textile production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Polymer dispersions/emulsions intended for use in paints and coatings, glues and adhesives, building materials and fluids used in textile production. Aim of application is the preservation of polymer mixtures which are consecutively used for formulation of paints, glues and further matrices (covered by PT 6).
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 26,4 - 107,1 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.8.1. *Use-specific instructions for use*

See general directions for use.

4.8.2. *Use-specific risk mitigation measures*

See general directions for use.

4.8.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.8.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.8.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.9. **Use description**

Table 9

In-can preservation of mineral slurries (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: -
Field(s) of use	indoor use Mineral slurries, e.g. CaCO ₃ -slurries as used for instance in paper industry and other industrial branches.
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 10,6 - 107,1 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 1,48 to 15 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

- 4.9.1.

Use-specific instructions for use

See general directions for use.
- 4.9.2.

Use-specific risk mitigation measures

See general directions for use.
- 4.9.3.

Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-
- 4.9.4.

Where specific to the use, the instructions for safe disposal of the product and its packaging

-
- 4.9.5.

Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-
- 4.10.

Use description

Table 10

Preservation of open re-circulating systems (PT11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in open recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: Dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.

Application rate(s) and frequency	Application Rate: a. Preventive: 5,3 – 107,1 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 31,4 - 107,1 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof shall be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast: 24 hours, against algae: 48 hours.
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.10.1. Use-specific instructions for use

See general directions for use.

4.10.2. Use-specific risk mitigation measures

The use is restricted to small cooling systems with a maximum blowdown of 2 m³/h. Waste water must be discharged to the municipal sewer or purified in an on-site industrial sewage treatment plant including a biological treatment step. The product can only be applied when the cooling towers are equipped with drift eliminators that reduce drift with at least 99 %.

See general directions for use.

4.10.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.10.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.10.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.11. Use description

Table 11

Preservation of cooling fluids in closed systems (PT11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids closed recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: Dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 5,3 – 107,1 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 31,4 - 107,1 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof shall be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast 24 hours, against algae 48 hours.

Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.11.1. *Use-specific instructions for use*

See general directions for use.

4.11.2. *Use-specific risk mitigation measures*

See general directions for use.

4.11.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.11.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.11.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.12. **Use description**

Table 12

Preservation of other liquids in e.g. air conditioning systems, air washers and pasteurizers (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). To protect water and other liquids in air conditioning systems, air washers, humidifiers, evaporate condensers, stationary sprinklers, water spray extinguishing systems and pasteurizers from microbial growth.

Application method(s)	Method: Dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 5,3 – 107,1 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm for preventive treatment against bacteria (incl. legionella), fungi and yeast, for preventive treatment against algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 31,4 - 107,1 mg product/kg matrix to be preserved. For curative treatment a pure CMIT/MIT concentration of 4,4 to 15 ppm shall be used. Number and timing of application: Frequency: Single or multiple application (number of additions depends on the state of the unit being treated) Contact time for curative use against bacteria (incl. legionella), fungi and yeast: 24 hours, against algae: 48 hours.
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

- 4.12.1.

Use-specific instructions for use

See general directions for use.
- 4.12.2.

Use-specific risk mitigation measures

See general directions for use.
- 4.12.3.

Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-
- 4.12.4.

Where specific to the use, the instructions for safe disposal of the product and its packaging

-
- 4.12.5.

Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-
- 4.13.

Use description

Table 13

Preservation of membrane module solutions (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: -
Field(s) of use	indoor use Membrane module solutions. Conservation / Preservation of the water in membrane units used in water pre-treatment preparation during production shutdown after cleaning. The biocides are applied for preservation of process fluids used for non-food membrane units/systems (e.g. reverse osmosis and ultrafiltration membranes), which are widely used in the water pre-treatment preparation (non-food, non potable water, non-medical). Membrane units are used in different industries (wastewater, surface technologies, sea water desalination, etc.). These systems are recirculating.
Application method(s)	Method: Dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: 73,6 – 121,4 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 10,3 to 17 ppm. Number and timing of application: Frequency: Single or multiple application (number of additions depends on the state of the unit being treated)
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.1.3.1. Use-specific instructions for use

During the shutdown of production, it is recommended to counteract microbiological growth inside the membrane elements. Before the shutdown, the plant has to be cleaned first, to remove all sorts of deposits like carbonates, salts, silica, organic matter or biomass. For this purpose, specialized membrane cleaners are used. After cleaning, the plant is flushed with permeate quality water until neutral pH is achieved. Only after these cleaning steps, CMIT/MIT containing water shall reach the plant and shall be regularly circulated by slow pumping. The biocidal product shall be added to the fill-water (permeate quality), to avoid microbiological growth during longer shut downs. During longer shut downs, the fill-water shall be regularly circulated by slow pumping and sampled to check for a microbiological re-contamination. A shift in pH can be a first indication of a re-contamination by microbiology. In this case the fill-up solution must be replaced.

See general directions for use.

4.13.2. Use-specific risk mitigation measures

See general directions for use.

4.13.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.13.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.13.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.14. Use description

Table 14

Preservation of white water circuits (short circulation) in paper industry (PT 12)

Product type	PT12: Slimicides
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Preservation of white water circuits (short circulation) in paper industry. a. To protect white water circuits (short circulation) in paper machines (preventive) against bacteria. b. For curative treatment in of white water circuits (short circulation) in paper machines already affected against bacteria and fungi. Aim of the application is to prevent or combat slime formation on pipework, and surfaces of materials and equipment in the paper area.
Application method(s)	Method: Dosing directly into the system (Shock / Continuous dosing). Detailed description: The biocidal product is dosed directly into the system to be protected. (Shock / Continuous dosing). Solutions of the biocidal product are automatically added by means of a dosing pump and pipes directly in the circuit (head box, mixing chest, collecting sump, broke, etc.), in intermittent or shock dosage.

Application rate(s) and frequency	Application Rate: a. Preventive: 5,3 – 107,1 mg product/kg matrix to be preserved (strongly dependent on actual system being treated). For preventive treatment (maintenance) in white water circuits (short circulation), the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,74 to 15 ppm. This refers to max. 10,6 g a.i./ton paper. b. Curative: 52,5 - 107,1 mg product/kg matrix to be preserved (strongly dependent on actual system being treated) For curative treatment (shock dose) in white water circuits (short circulation), the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 7,35 – 15 ppm. Number and timing of application: Frequency: The number of additions per week depends on the state of the unit being treated. The addition frequency of biocidal product in shock dosage is 1 to 6 times per day. Contact time for curative use: 24 hours.
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.14.1. Use-specific instructions for use

See general directions for use.

4.14.2. Use-specific risk mitigation measures

Application is only allowed in paper factories that comply with the Industrial Emission Directive 2010/75/EU where wastewater is purified in an on-site industrial sewage treatment plant including a biological treatment step in accordance to the Best Available Techniques (BAT) as prescribed in the BAT-reference document (BREF) for the production of pulp, paper and board. The effluent must be diluted at least 200 times. Paper factories that are exempted from the Industrial Emission Directive must discharge to the municipal sewer.

See general directions for use.

4.14.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.14.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.14.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.15. Use description

Table 15

Preservation of liquids used to treat or cut metal and glass (PT 13)

Product type	PT13: Working or cutting fluid preservatives
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use For use as a protective agent in liquids used to treat or cut metal and glass. Aim of application is to prevent microbial deterioration caused by bacteria, fungi or yeasts.
Application method(s)	Method: Dosing directly into the system. Detailed description: The biocidal product is added to the emulsion already formed. The biocide is added mechanical or fully automated with a timer. For tank-side addition, the biocidal products are added with an automated dosing pump and specific pipes into the collecting sump where the metal working fluids flow. The product introduction needs to be done below the water level to enable a fast mixing and reduce the biocide exposure risk. The dosage is done for 30-60 minutes.
Application rate(s) and frequency	Application Rate: 169,3 – 214,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 23,7 to 30 ppm. Number and timing of application: Frequency: 1-7 times a week.
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.15.1. *Use-specific instructions for use*

See general directions for use.

4.15.2. *Use-specific risk mitigation measures*

See general directions for use.

4.15.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.15.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.15.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

5. GENERAL DIRECTIONS FOR USE OF THE META SPC 1

5.1. Instructions for use

General instructions

Preliminary tests to determine suitability, compatibility and optimal concentration are recommended prior to the use of the product in a new application. For more technical information, see the currently valid product information sheet which will be provided to customers upon request.

The product labels give instructions on the dosage of the product.

General use information

PT 6

For the in-can preservation of washing and cleaning fluids, paints and coatings, fluids used in paper, textile and leather production, as well as glues and adhesives, concrete additives and building materials (e.g. fillers/sealants, plasters, wax emulsions), polymer dispersions/emulsions (intended for use in paints, coatings, glues, adhesives, building materials and fluids used in textile production) and the treatment of mineral slurries.

The product must be homogeneously incorporated into the matrix to be protected.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

PT 11

For use as a biocide for the preservation of cooling fluids in circulating systems and other liquids in e.g. air conditioning systems, air washers and pasteurizers from infestation with harmful organisms such as bacteria (including Legionella), fungi, yeasts and algae. Furthermore, the applications covered by PT 11 comprise the preservation of membrane module solutions.

Heavily contaminated units shall be cleaned prior to treatment.

The biocidal product shall be added to the water circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The microbiocidal effect begins immediately after metering.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used. Due to variation in the different systems and matrices (organic load, microbial contamination, slime forming, temperature, pH, etc.) accurate measurements by chemical and microbiological tests shall be carried out to determine the effective dose for the specific location or system.

PT 12

For use as a biocide for the preservation of white water circuits (short circulation) in paper machines. Aim of the application is to prevent or combat slime formation (as caused by bacteria and fungi) on pipework and surfaces of materials and equipment in the paper area.

The microbiocidal effect begins immediately after metering.

The biocidal product shall be added to the white water circuit (short circulation) in paper machines to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

PT 13

For use as a protective agent for the preservation of liquids used to treat or cut metal and glass. Aim of application is to prevent microbial deterioration caused by bacteria, fungi or yeasts.

The ready-to-use biocidal product, without former dilution or formulation, is directly applied into the ready to use metalworking fluid cycle to control microbial deterioration and to maintain the correct function of the metalworking fluid.

The microbiocidal effect begins immediately after metering.

The biocidal product is added to the metalworking circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

5.2. Risk mitigation measures

During product handling: Wear protective gloves consistent with European Standard EN ISO 374 or equivalent, coverall (Chemical protection clothes type 6) consistent with European Standard EN ISO 13034 or equivalent and eye or face protection (recommended splash goggles or face shield) consistent with European Standard EN ISO 16321 or equivalent.

In case of C(M)IT/MIT concentrations ≥ 15 ppm: During product handling, application and post-application: Wear protective chemical resistant gloves consistent with European Standard EN ISO 374 or equivalent and a protective coverall (at least type 6) consistent with European Standard EN ISO 13034 or equivalent which is impermeable for the biocidal product.

Equipment should preferably be cleaned with water. Dilute rinsing water sufficiently before disposal via a wastewater treatment plant.

Only vent waste air to the atmosphere via suitable separators or scrubbers. No special fire or explosion protection measures necessary.

Required Personal Protection Equipment is further described in Safety Data Sheet.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*First aid measures*

IF SWALLOWED: Immediately rinse mouth. Give something to drink, if exposed person is able to swallow. Do NOT induce vomiting. Call 112/ambulance for medical assistance.

Information to Healthcare personnel/doctor:

Initiate life support measures if needed, thereafter call a POISON CENTRE.

IF INHALED: Move to fresh air and keep at rest in a position comfortable for breathing.

If symptoms: Call 112/ambulance for medical assistance.

If no symptoms: Call a POISON CENTRE or a doctor.

Information to Healthcare personnel/doctor:

Initiate life support measures if needed, thereafter call a POISON CENTRE.

IF ON SKIN: Immediately wash skin with plenty of water. Thereafter take off all contaminated clothing and wash it before reuse. Continue to wash the skin with water and continue rinsing for 15 minutes. Call a POISON CENTRE or a doctor.

Information to Healthcare personnel/doctor:

Initiate life support measures if needed, thereafter call a POISON CENTRE.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing for at least 15 minutes. Call 112/ambulance for medical assistance.

Particular environmental risks

Product containers shall be carefully handled, stored and transported to prevent damage to the containers and leakage of the product into soil, air and water. The biocidal product is harmful to aquatic organisms. The product is biodegradable following dilution.

5.4. Instructions for safe disposal of the product and its packaging

Disposal information

Check whether re-use of the product in the packaging is possible. Pack or seal product waste and contaminated empty containers, label, and discharge in accordance with refuse disposal acts.

In the case of larger amounts of product in the packaging, discuss with the supplier of the product. The recipient of contaminated empty containers must be informed of the possible risks due to product residues. In the case of disposal within the EU, use the respective valid waste code in accordance with the European Waste Catalogue (EWC). The producer of the waste is responsible for assigning sector- and process-specific waste codes to his waste in accordance with the European Waste Catalogue.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Shelf-life: 24 months.

Store in tightly sealed original containers. Protect from frost.

6. OTHER INFORMATION

With respect to the "Category (ies) of users" note: "Professionals (including industrial users) means trained professionals if this is required by national legislation."

Resistance management for intended applications (PT 11, 12, 13):

- avoidance of under-dosing;
- frequent detection of efficacy and of biocide content in the industrial systems to assure maintenance of correct CMIT/MIT concentration;
- for new applications, preliminary tests to determine the suitability, compatibility and optimal application concentration are strongly recommended;
- under difficult conditions alternation of active ingredients, i.e. rotation with other biocides, also combination with other products may be useful.

Full titles of EN standards and legislation referred to in section 5.2:

EN ISO 374 – Protective gloves against dangerous chemicals and micro-organisms.

EN ISO 13034 - Protective clothing against liquid chemicals - Performance requirements for chemical protective clothing offering limited protective performance against liquid chemicals (Type 6 and Type PB [6] equipment)

EN ISO 16321 – Eye and face protection for occupational use

7. THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 1

7.1. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preventol IT14	Market area: EU		
			Kathon® CMI 14	Market area: EU		
Authorisation number			EU-0031652-0001 1-1			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		24,2 % (w/w)	

7.2. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preventol IT14 MV	Market area: EU		
			Kathon® CMI 14 MV	Market area: EU		
Authorisation number			EU-0031652-0002 1-1			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol-3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		24,2 % (w/w)	

7.3. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Kurita F-5000	Market area: EU		
Authorisation number			EU-0031652-0003 1-1			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		24,2 % (w/w)	

7.4. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 01-04	Market area: EU		
Authorisation number			EU-0031652-0004 1-1			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol-3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		24,2 % (w/w)	

7.5. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 01-05	Market area: EU		
Authorisation number			EU-0031652-0005 1-1			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol-3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		24,2 % (w/w)	

1. META SPC 2 ADMINISTRATIVE INFORMATION

1.1. Meta SPC 2 identifier

Identifier	Meta SPC: Meta SPC 2
------------	----------------------

1.2. Suffix to the authorisation number

Number	1-2
--------	-----

1.3. Product type(s)

Product type(s)	PT11: Preservatives for liquid-cooling and processing systems
-----------------	---

2. META SPC 2 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 2

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		5 - 5,51 % (w/w)

2.2. Type(s) of formulation of the meta SPC 2

Formulation type(s)	AL Any other liquid
---------------------	---------------------

3. HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 2

Hazard statements	H290: May be corrosive to metals. H314: Causes severe skin burns and eye damage. H317: May cause an allergic skin reaction. H410: Very toxic to aquatic life with long lasting effects. EUH071: Corrosive to the respiratory tract.
Precautionary statements	P234: Keep only in original packaging. P272: Contaminated work clothing should not be allowed out of the workplace. P273: Avoid release to the environment. P280: Wear protective gloves. P280: Wear protective clothing. P280: Wear eye protection. P280: Wear face protection. P301+P330+P331: IF SWALLOWED: rinse mouth. Do NOT induce vomiting. P310: Immediately call a doctor. P310: Immediately call a POISON CENTER. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water [or shower]. P310: Immediately call a doctor. P310: Immediately call a POISON CENTER. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P310: Immediately call a doctor. P310: Immediately call a POISON CENTER. P362+P364: Take off contaminated clothing and wash it before reuse. P390: Absorb spillage to prevent material damage. P405: Store locked up. P406: Store in a corrosive-resistant container with a resistant inner liner. P501: Dispose of contents in accordance with all local, regional, national and international regulations. P501: Dispose of container in accordance with all local, regional, national and international regulations. P391: Collect spillage.

4. AUTHORISED USE(S) OF THE META SPC

4.1. Use description

Table 1

Preservation of cooling fluids in open recirculating systems (PT11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in open recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: Dosing directly into the system. Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 23,0 – 517,2 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 137,9 - 517,2 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof shall be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast: 24 hours, against algae: 48 hours.

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.1.1. *Use-specific instructions for use*

See general directions for use.

4.1.2. *Use-specific risk mitigation measures*

The use is restricted to small cooling systems with a maximum blowdown of 2 m³/h. Waste water shall be discharged to the municipal sewer or purified in an on-site industrial sewage treatment plant including a biological treatment step. The product shall only be applied when the cooling towers are equipped with drift eliminators that reduce drift with at least 99 %.

See general directions for use.

4.1.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.1.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.1.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.2. **Use description**

Table 2

Preservation of cooling fluids in closed systems (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -

Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in closed recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: Dosing directly into the system. Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 23,0 – 517,2 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into matrix to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 137,9 - 517,2 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof shall be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast: 24 hours, against algae: 48 hours.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.2.1. *Use-specific instructions for use*

See general directions for use.

4.2.2. *Use-specific risk mitigation measures*

See general directions for use.

4.2.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.2.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.2.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

5. GENERAL DIRECTIONS FOR USE OF THE META SPC 2

5.1. Instructions for use

General instructions

Preliminary tests to determine suitability, compatibility and optimal concentration are recommended prior to the use of the product in a new application. For more technical information, see the currently valid product information sheet which will be provided to customers upon request.

The product labels give instructions on the dosage of the product.

General use information

PT 11

For use as a biocide for the preservation of cooling fluids in circulating systems from infestation with harmful organisms such as bacteria (including Legionella), fungi, yeasts and algae.

Heavily infested units shall be cleaned prior to treatment.

The biocidal product is added to the water circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The microbiocidal effect begins immediately after metering.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

5.2. Risk mitigation measures

During product handling: Wear protective gloves consistent with European Standard EN ISO 374 or equivalent, coverall (Chemical protection clothes type 6) consistent with European Standard EN ISO 13034 or equivalent and eye or face protection (recommended splash goggles or face shield) consistent with European Standard EN ISO 16321 or equivalent.

In case of C(M)IT/MIT concentrations ≥ 15 ppm: During product handling, application and post-application: Wear protective chemical resistant gloves consistent with European Standard EN ISO 374 or equivalent and a protective coverall (at least type 6) consistent with European Standard EN ISO 13034 or equivalent which is impermeable for the biocidal product.

Equipment should preferably be cleaned with water. Dilute rinsing water sufficiently before disposal via a wastewater treatment plant.

Only vent waste air to the atmosphere via suitable separators or scrubbers. No special fire or explosion protection measures necessary.

Required Personal Protection Equipment is described in Safety Data Sheet.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

First aid measures

IF SWALLOWED: Immediately rinse mouth. Give something to drink, if exposed person is able to swallow. Do NOT induce vomiting. Call 112/ambulance for medical assistance.

IF INHALED: Move to fresh air and keep at rest in a position comfortable for breathing.

If symptoms: Call 112/ambulance for medical assistance.

If no symptoms: Call a POISON CENTRE or a doctor.

IF ON SKIN: Immediately wash skin with plenty of water. Thereafter take off all contaminated clothing and wash it before reuse. Continue to wash the skin with water for 15 minutes. Call a POISON CENTRE or a doctor.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing for at least 15 minutes. Call 112/ambulance for medical assistance.

Particular environmental risks

Product containers shall be carefully handled, stored and transported to prevent damage to the containers and leakage of the product into soil, air and water. The biocidal product is harmful to aquatic organisms. The product is biodegradable following dilution.

5.4. **Instructions for safe disposal of the product and its packaging**

Disposal information

Check whether re-use of the product in the packaging is possible. Pack or seal product waste and contaminated empty containers, label, and discharge in accordance with refuse disposal acts.

In the case of larger amounts of product in the packaging, discuss with the supplier of the product. The recipient of contaminated empty containers shall be informed of the possible risks due to product residues. In the case of disposal within the EU, use the respective valid waste code in accordance with the European Waste Catalogue (EWC). The producer of the waste is responsible for assigning sector- and process-specific waste codes to his waste in accordance with the European Waste Catalogue.

5.5. **Conditions of storage and shelf-life of the product under normal conditions of storage**

Shelf-life: 24 months.

Store in tightly sealed original containers. Protect from frost.

6. **OTHER INFORMATION**

With respect to the “Category (ies) of users” note: “Professionals (including industrial users) means trained professionals if this is required by national legislation.”

Resistance management for intended applications (PT 11):

- avoidance of under-dosing;
- frequent detection of efficacy and of biocide content in the industrial systems to assure maintenance of correct CMIT/MIT concentration;
- for new applications, preliminary tests to determine the suitability, compatibility and optimal application concentration are strongly recommended;
- under difficult conditions alternation of active ingredients, i.e. rotation with other biocides, also combination with other products may be useful.

Full titles of EN standards and legislation referred to in section 5.2:

EN ISO 374 – Protective gloves against dangerous chemicals and micro-organisms.

EN ISO 13034 – Protective clothing against liquid chemicals - Performance requirements for chemical protective clothing offering limited protective performance against liquid chemicals (Type 6 and Type PB [6] equipment)

EN ISO 16321 – Eye and face protection for occupational use

7. THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 2

7.1. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 02-01	Market area: EU		
Authorisation number			EU-0031652-0006 1-2			
Common name	IUPAC name	Function	CAS number	EC number		Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9			5,2 % (w/w)

7.2. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 02-02	Market area: EU	
Authorisation number			EU-0031652-0007 1-2		
Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol-3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		5,2 % (w/w)

1. META SPC 3 ADMINISTRATIVE INFORMATION

1.1. Meta SPC 3 identifier

Identifier	Meta SPC: Meta SPC 3
------------	----------------------

1.2. Suffix to the authorisation number

Number	1-3
--------	-----

1.3. Product type(s)

Product type(s)	PT06: Preservatives for products during storage PT11: Preservatives for liquid-cooling and processing systems PT12: Slimicides PT13: Working or cutting fluid preservatives
-----------------	---

2. META SPC 3 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 3

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		1,7 - 2,8 % (w/w)

2.2. Type(s) of formulation of the meta SPC 3

Formulation type(s)	AL Any other liquid
---------------------	---------------------

3. HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 3

Hazard statements	H314: Causes severe skin burns and eye damage. H317: May cause an allergic skin reaction. H410: Very toxic to aquatic life with long lasting effects. EUH071: Corrosive to the respiratory tract.
Precautionary statements	P272: Contaminated work clothing should not be allowed out of the workplace. P273: Avoid release to the environment. P280: Wear protective gloves. P280: Wear protective clothing. P280: Wear eye protection. P280: Wear face protection. P301+P330+P331: IF SWALLOWED: rinse mouth. Do NOT induce vomiting. P310: Immediately call a doctor. P310: Immediately call a POISON CENTER. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water [or shower]. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P362+P364: Take off contaminated clothing and wash it before reuse. P405: Store locked up. P501: Dispose of contents in accordance with all local, regional, national and international regulations.. P501: Dispose of container in accordance with all local, regional, national and international regulations.. P391: Collect spillage.

4. AUTHORISED USE(S) OF THE META SPC

4.1. Use description

Table 1

In-can preservation of washing and cleaning fluids (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Washing and cleaning fluids
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected during the production. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 231,3 – 1500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected in such a way that a pure CMIT/MIT concentration of 3,7 to less than 15 ppm is maintained. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.1.1. Use-specific instructions for use

See general directions for use.

4.1.2. Use-specific risk mitigation measures

See general directions for use.

4.1.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.1.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.1.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.2. Use description

Table 2

In-can preservation of paints and coatings (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Paints and coatings
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 695,7 – 1 500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 11,1 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.2.1. *Use-specific instructions for use*

See general directions for use.

4.2.2. *Use-specific risk mitigation measures*

See general directions for use.

4.2.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.2.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.2.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.3. **Use description**

Table 3

In-can preservation of fluids used in paper production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in paper production (e.g. pigment slurries)
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 462,5 – 3000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.3.1. *Use-specific instructions for use*

See general directions for use.

4.3.2. *Use-specific risk mitigation measures*

See general directions for use.

4.3.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.3.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.3.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.4. **Use description**

Table 4

In-can preservation of fluids used in textile production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in textile production
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 231,3 – 3000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.4.1. *Use-specific instructions for use*

See general directions for use.

4.4.2. Use-specific risk mitigation measures

See general directions for use.

4.4.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.4.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.4.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.5. Use description

Table 5

In-can preservation of fluids used in leather production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in leather production
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected.The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 462,5 – 3000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 30 ppm. Number and timing of application: Frequency: Single application

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.5.1. *Use-specific instructions for use*

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

See general directions for use.

4.5.2. *Use-specific risk mitigation measures*

See general directions for use.

4.5.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.5.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.5.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.6. **Use description**

Table 6

In-can preservation of glues and adhesives (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: Common name: fungi Development stage: -
Field(s) of use	indoor use Glues and adhesives
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.

Application rate(s) and frequency	Application Rate: 231,3 – 3000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 30 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.6.1. *Use-specific instructions for use*

See general directions for use.

4.6.2. *Use-specific risk mitigation measures*

See general directions for use.

4.6.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.6.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.6.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.7. **Use description**

Table 7

In-can preservation of concrete additives and building material such as fillers/sealants, plasters and wax emulsions (PT 6) for indoor use

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -

Field(s) of use	indoor use Concrete additives and building material such as fillers/sealants, plasters and wax emulsions intended for indoor use.
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected during the production. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 462,5 - 1500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 15 ppm. End products which are used by the general public must contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.7.1. *Use-specific instructions for use*

See general directions for use.

4.7.2. *Use-specific risk mitigation measures*

See general directions for use.

4.7.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.7.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.7.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.8. **Use description**

Table 8

In-can preservation of polymer dispersions/emulsions intended for use in paints and coatings, glues and adhesives, building materials and fluids used in textile production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Polymer dispersion/emulsions intended for use in paints and coatings, glues and adhesives, building materials and fluids used in textile production. Aim of application is the preservation of polymer mixtures which are consecutively used for formulation of paints, glues and further matrices (covered by PT 6).
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 231,3 - 1500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.8.1. *Use-specific instructions for use*

See general directions for use.

4.8.2. *Use-specific risk mitigation measures*

See general directions for use.

4.8.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.8.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.8.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.9. Use description

Table 9

In-can preservation of mineral slurries (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: -
Field(s) of use	indoor use Mineral slurries, e.g. CaCO ₃ -slurries as used for instance in paper industry and other industrial branches.
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 92,5 - 1 500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 1,48 to 15 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.9.1. Use-specific instructions for use

See general directions for use.

4.9.2. Use-specific risk mitigation measures

See general directions for use.

4.9.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.9.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.9.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.10. Use description

Table 10

Preservation of cooling fluids in open circulating systems (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in open recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 45,9 – 1500 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 275,0 - 1500 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof shall be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast 24 hours, against algae 48 hours.

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.10.1. *Use-specific instructions for use*

See general directions for use.

4.10.2. *Use-specific risk mitigation measures*

The use is restricted to small cooling systems with a maximum blowdown of 2 m³/h. Waste water must be discharged to the municipal sewer or purified in an on-site industrial sewage treatment plant including a biological treatment step. The product shall only be applied when the cooling towers are equipped with drift eliminators that reduce drift with at least 99 %.

See general directions for use.

4.10.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.10.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.10.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.11. **Use description**

Table 11

Preservation of cooling fluids in closed systems (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -

Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in closed recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 45,9 – 1500 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 0.735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1.42 to 15 ppm shall be used. b. Curative: 275,0 - 1500 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof shall be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast: 24 hours, against algae: 48 hours.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.11.1. *Use-specific instructions for use*

See general directions for use.

4.11.2. *Use-specific risk mitigation measures*

See general directions for use.

4.11.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.11.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.11.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.12. Use description

Table 12

Preservation of other liquids in e.g. air conditioning systems, air washers and pasteurizers (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). To protect water and other liquids in air conditioning systems, air washers, humidifiers, evaporate condensers, stationary sprinklers, water spray extinguishing systems and pasteurizers, from microbial growth.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 45,9 – 1500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into liquids used in air conditioning systems, air washers, humidifiers, evaporate condensers, membrane systems, stationary sprinkler, water spray extinguishing systems and pasteurizers to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm for preventive treatment against bacteria (incl. legionella), fungi and yeast, for preventive treatment against algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 275,0 - 1500 mg product/kg matrix to be preserved. For curative treatment a pure CMIT/MIT concentration of 4,4 to 15 ppm shall be used. Number and timing of application: Frequency: Single or multiple application (number of additions depends on the state of the unit being treated). Contact time for curative use against bacteria (incl. legionella), fungi and yeast: 24 hours, against algae: 48 hours.
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.12.1. *Use-specific instructions for use*

See general directions for use.

4.12.2. *Use-specific risk mitigation measures*

See general directions for use.

4.12.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.12.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.12.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.13. **Use description**

Table 13

Preservation of wood preservative treatment solutions (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Preservation of wood preservative treatment solutions
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the wood preservative solution to be protected. Solutions of the biocidal product are automatically dosed in the wood preservative solution during the flow.
Application rate(s) and frequency	Application Rate: 2869 to 5000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into wood preservative treatment solution to be protected leading to a pure CMIT/MIT concentration of 45,9 to 50 ppm. Number and timing of application: Frequency: Number of additions per week depends on the state of the unit being treated.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.13.1. *Use-specific instructions for use*

See general directions for use.

4.13.2. Use-specific risk mitigation measures

The product shall not be applied to wood preservative treatment solutions which are used to preserve wood intended for direct contact with water or soils (use class 4).

Freshly treated timber must be stored after treatment under shelter or on impermeable hard standing, or both, to prevent direct losses to soil, sewer or water and that any losses shall be collected for reuse or disposal.

See general directions for use.

4.13.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.13.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.13.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.14. Use description

Table 14

Preservation of membrane module solutions (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: -
Field(s) of use	indoor use Membrane module solutions. Conservation / Preservation of the water in membrane units used in water pre-treatment preparation during production shutdown after cleaning. The biocides are applied for preservation of process fluids used for non-food membrane units/systems (e.g. reverse osmosis and ultrafiltration membranes), which are widely used in the water pre-treatment preparation (non-food, non potable water, non-medical). Membrane units are used in different industries (wastewater, surface technologies, sea water desalination, etc.). These systems are recirculating.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.

Application rate(s) and frequency	Application Rate: 643,8 – 1700 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 10,3 to 17 ppm. Number and timing of application: Frequency: Single or multiple application (number of additions depends on the state of the unit being treated)
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.14.1. *Use-specific instructions for use*

During the shutdown of production, it is recommended to counteract microbiological growth inside the membrane elements. Before the shutdown, the plant shall be cleaned first, to remove all sorts of deposits like carbonates, salts, silica, organic matter or biomass. For this purpose, specialized membrane cleaners are used. After cleaning, the plant is flushed with permeate quality water until neutral pH is achieved. Only after these cleaning steps, CMIT/MIT containing water is reaching the plant and is regularly circulated by slow pumping. The biocidal product shall be added to the fill-water (permeate quality), to avoid microbiological growth during longer shutdowns. During longer shutdowns, the fill-water shall be regularly circulated by slow pumping and sampled to check for a microbiological re-contamination. A shift in pH can be a first indication of a re-contamination by microbiology. In this case the fill-up solution shall be replaced.

See general directions for use.

4.14.2. *Use-specific risk mitigation measures*

See general directions for use.

4.14.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.14.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.14.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.15. **Use description**

Table 15

Preservation of white water circuits (short circulation) in paper industry (PT 12)

Product type	PT12: Slimicides
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Preservation of white water circuits (short circulation) in paper industry. a. To protect white water circuits (short circulation) in paper machines (preventive) against bacteria. b. For curative treatment in of white water circuits (short circulation) in paper machines already affected against bacteria and fungi. Aim of the application is to prevent or combat slime formation on pipework, and surfaces of materials and equipment in the paper area.
Application method(s)	Method: dosing directly into the system (Shock / Continuous dosing) Detailed description: The biocidal product is dosed directly into the system to be protected. (Shock / Continuous dosing). Solutions of the biocidal product are automatically added by means of a dosing pump and pipes directly in the circuit (head box, mixing chest, collecting sump, broke, etc.), in intermittent or shock dosage.
Application rate(s) and frequency	Application Rate: a. Preventive: 46,3 – 1 500 mg product/kg matrix to be preserved (strongly dependent on actual system being treated). For preventive treatment (maintenance) in white water circuits (short circulation), the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,74 to 15 ppm. This refers to max. 10,6 g a.i./ton paper. b. Curative: 459,4 – 1 500 mg product/kg matrix to be preserved (strongly dependent on actual system being treated). For curative treatment (shock dose) in white water circuits (short circulation), the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 7,35 to 15 ppm. Number and timing of application: Frequency: The number of additions per week depends on the state of the unit being treated. The addition frequency of biocidal product in shock dosage is 1 to 6 times per day. Contact time for curative use: 24 hours
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.15.1. Use-specific instructions for use

See general directions for use.

4.15.2. *Use-specific risk mitigation measures*

Application is only allowed in paper factories that comply with the Industrial Emission Directive 2010/75/EU where wastewater is purified in an on-site industrial sewage treatment plant including a biological treatment step in accordance to the Best Available Techniques (BAT) as prescribed in the BAT-reference document (BREF) for the production of pulp, paper and board. The effluent must be diluted at least 200 times. Paper factories that are exempted from the Industrial Emission Directive shall discharge to the municipal sewer.

See general directions for use.

4.15.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.15.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.15.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.16. **Use description**

Table 16

Preservation of liquids used to treat or cut metal and glass (PT 13)

Product type	PT13: Working or cutting fluid preservatives
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use For use as a protective agent in liquids used to treat or cut metal and glass. Aim of application is to prevent microbial deterioration caused by bacteria, fungi or yeasts.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is added to the emulsion already formed. The biocide is added mechanical or fully automated with a timer. For tank-side addition, the biocidal products are added with an automated dosing pump and specific pipes into the collecting sump where the metal working fluids flow. The product introduction needs to be done below the water level to enable a fast mixing and reduce the biocide exposure risk. The dosage is done for 30-60 minutes.

Application rate(s) and frequency	Application Rate: 1 481,3 - 3 000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 23,7 to 30 ppm. Number and timing of application: Frequency: 1-7 times a week
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.16.1. *Use-specific instructions for use*

See general directions for use.

4.16.2. *Use-specific risk mitigation measures*

See general directions for use.

4.16.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.16.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.16.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

5. **GENERAL DIRECTIONS FOR USE OF THE META SPC 3**

5.1. **Instructions for use**

General instructions

Preliminary tests to determine suitability, compatibility and optimal concentration are recommended prior to the use of the product in a new application. For more technical information, see the currently valid product information sheet which will be provided to the customer on request.

The product labels give instructions on the dosage of the product.

General use information

PT 6

For the in-can preservation of washing and cleaning fluids, paints and coatings, fluids used in paper, textile and leather production, as well as glues and adhesives, concrete additives and building materials (e.g. fillers/sealants, plasters, wax emulsions), polymer dispersions/ emulsions (intended for use in paints, coatings, glues, adhesives, building materials and fluids used in textile production) and the treatment of mineral slurries.

The product must be homogeneously incorporated into the matrix to be protected.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

PT 11

For use as a biocide for the preservation of cooling fluids in circulating systems and other liquids in e.g. air conditioning systems, air washers and pasteurizers from infestation with harmful organisms such as bacteria (including *Legionella*), fungi, yeasts and algae. Furthermore, the applications covered by PT 11 comprise the preservation of membrane module solutions and wood preservative treatment solutions.

Heavily contaminated units shall be cleaned prior to treatment.

The biocidal product is added to the water circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The microbiocidal effect begins immediately after metering.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used. Due to variation in the different systems and matrices (organic load, microbial contamination, slime forming, temperature, pH, etc.) accurate measurements by chemical and microbiological tests shall be carried out to determine the effective dose for the specific location or system.

PT 12

For use as a biocide to for the preservation of white water circuits (short circulation) in paper machines. Aim of the application is to prevent or combat slime formation (as caused by bacteria and fungi) on pipework and surfaces of materials and equipment in the paper area.

The microbiocidal effect begins immediately after metering.

The biocidal product is added to the white water circuit (short circulation) in paper machines to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

PT 13

For use as a protective agent for the preservation of liquids used to treat or cut metal and glass. Aim of application is to prevent microbial deterioration caused by bacteria, fungi or yeasts.

A ready-to-use biocidal product, without former dilution or formulation), is directly applied into the ready to use metalworking fluid cycle to control microbial deterioration and to maintain the correct function of the metalworking fluid.

The microbiocidal effect begins immediately after metering.

The biocidal product is added to the metalworking circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

5.2. Risk mitigation measures

During product handling: Wear protective gloves consistent with European Standard EN ISO 374 or equivalent, overall (Chemical protection clothes type 6) consistent with European Standard EN ISO EN 13034 or equivalent and eye or face protection (recommended splash goggles or face shield) consistent with European Standard EN ISO 16321 or equivalent.

In case of C(M)IT/MIT concentrations ≥ 15 ppm: During product handling, application and post-application: Wear protective chemical resistant gloves consistent with European Standard EN ISO 374 or equivalent and a protective coverall (at least type 6) consistent with European Standard EN ISO 13034 or equivalent which is impermeable for the biocidal product.

Equipment should preferably be cleaned with water. Dilute rinsing water sufficiently before disposal via a wastewater treatment plant.

Only vent waste air to the atmosphere via suitable separators or scrubbers. No special fire or explosion protection measures necessary.

Required Personal Protection Equipment is described in Safety Data Sheet.

5.3. **Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment**

First aid measures

IF SWALLOWED: Immediately rinse mouth. Give something to drink, if exposed person is able to swallow. Do NOT induce vomiting. Call 112/ambulance for medical assistance.

IF INHALED: Move to fresh air and keep at rest in a position comfortable for breathing.

If symptoms: Call 112/ambulance for medical assistance.

If no symptoms: Call a POISON CENTRE or a doctor.

IF ON SKIN: Immediately wash skin with plenty of water. Thereafter take off all contaminated clothing and wash it before reuse. Continue to wash the skin with water for 15 minutes. Call a POISON CENTRE or a doctor.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing for at least 15 minutes. Call 112/ambulance for medical assistance.

Particular environmental risks

Product containers shall be carefully handled, stored and transported to prevent damage to the containers and leakage of the product into soil, air and water. The biocidal product is harmful to aquatic organisms. The product is biodegradable following dilution.

5.4. **Instructions for safe disposal of the product and its packaging**

Disposal information

Check whether re-use of the product in the packaging is possible. Pack or seal product waste and contaminated empty containers, label, and discharge in accordance with refuse disposal acts.

In the case of larger amounts of product in the packaging, discuss with the supplier of the product. The recipient of contaminated empty containers must be informed of the possible risks due to product residues. In the case of disposal within the EU, use the respective valid waste code in accordance with the European Waste Catalogue (EWC). The producer of the waste is responsible for assigning sector- and process-specific waste codes to his waste in accordance with the European Waste Catalogue.

5.5. **Conditions of storage and shelf-life of the product under normal conditions of storage**

Shelf-life: 24 months.

Store in tightly sealed original containers. Protect from frost.

6. OTHER INFORMATION

With respect to the “Category (ies) of users” note: “Professionals (including industrial users) means trained professionals if this is required by national legislation.”

Resistance management for intended applications (PT 11, 12, 13):

- avoidance of under-dosing;
- frequent detection of efficacy and of biocide content in the industrial systems to assure maintenance of correct CMIT/MIT concentration;
- for new applications, preliminary tests to determine the suitability, compatibility and optimal application concentration are strongly recommended;
- under difficult conditions alternation of active ingredients, i.e. rotation with other biocides, also combination with other products may be useful.

Full titles of EN standards and legislation referred to in section 5.2:

EN ISO 374 – Protective gloves against dangerous chemicals and micro-organisms.

EN ISO 13034 - Protective clothing against liquid chemicals - Performance requirements for chemical protective clothing offering limited protective performance against liquid chemicals (Type 6 and Type PB [6] equipment)

EN ISO 16321 – Eye and face protection for occupational use

7. THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 3

7.1. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)	Preventol D7	Market area: EU			
	Kathon® CMI 1.5	Market area: EU			
	BW425	Market area: EU			
	CELLAQUA IS	Market area: EU			
	CELLOSMOSE	Market area: EU			
	TDF WATER 6008	Market area: EU			
	ANTRAIN 6260	Market area: EU			
	BIDAQUA 37	Market area: EU			
	CIRA B10	Market area: EU			
Authorisation number		EU-0031652-0008 1-3			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		2,7 % (w/w)

7.2. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preventol D7 LT	Market area: EU		
			Kathon® CMI 1.5 MV	Market area: EU		
			S-Care-IT20	Market area: EU		
			ALTO*107 L	Market area: EU		
			MIRECIDE-KW/52	Market area: EU		
Authorisation number			EU-0031652-0009 1-3			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		2,6 % (w/w)	

7.3. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 03-05	Market area: EU		
Authorisation number			EU-0031652-0010 1-3			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,4 % (w/w)	

7.4. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)	Preservative 03-06	Market area: EU
Authorisation number	EU-0031652-0011 1-3	

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,6 % (w/w)

7.5. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 03-07	Market area: EU	
Authorisation number			EU-0031652-0012 1-3		
Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,6 % (w/w)

7.6. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 03-09	Market area: EU		
Authorisation number			EU-0031652-0013 1-3			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,7 % (w/w)	

7.7. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 03-10	Market area: EU				
Authorisation number			EU-0031652-0014 1-3					
Common name	IUPAC name	Function		CAS number		EC number		Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance		55965-84-9				2,6 % (w/w)

7.8. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Wolsit SP	Market area: EU		
Authorisation number			EU-0031652-0015 1-3			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		2,7 % (w/w)	

1. META SPC 4 ADMINISTRATIVE INFORMATION

1.1. Meta SPC 4 identifier

Identifier	Meta SPC: Meta SPC 4
------------	----------------------

1.2. Suffix to the authorisation number

Number	1-4
--------	-----

1.3. Product type(s)

Product type(s)	PT06: Preservatives for products during storage PT11: Preservatives for liquid-cooling and processing systems PT12: Slimicides PT13: Working or cutting fluid preservatives
-----------------	--

2. META SPC 4 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 4

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		3,6 - 5 % (w/w)

2.2. Type(s) of formulation of the meta SPC 4

Formulation type(s)	AL Any other liquid
---------------------	---------------------

3. HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 4

Hazard statements	H290: May be corrosive to metals. H314: Causes severe skin burns and eye damage. H317: May cause an allergic skin reaction. H410: Very toxic to aquatic life with long lasting effects. EUH071: Corrosive to the respiratory tract.
Precautionary statements	P234: Keep only in original packaging. P272: Contaminated work clothing should not be allowed out of the workplace. P273: Avoid release to the environment. P280: Wear protective gloves. P280: Wear protective clothing. P280: Wear eye protection. P280: Wear face protection. P301+P330+P331: IF SWALLOWED: rinse mouth. Do NOT induce vomiting. P310: Immediately call a doctor. P310: Immediately call a POISON CENTER. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water [or shower]. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

	P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P362+P364: Take off contaminated clothing and wash it before reuse. P390: Absorb spillage to prevent material damage. P405: Store locked up. P406: Store in a corrosive-resistant container with a resistant inner liner. P501: Dispose of contents in accordance with all local, regional, national and international regulations.. P501: Dispose of container in accordance with all local, regional, national and international regulations.. P391: Collect spillage.
--	--

4. AUTHORISED USE(S) OF THE META SPC

4.1. Use description

Table 1

In-can preservation of washing and cleaning fluids (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Washing and cleaning fluids
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected during the production. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 127,6 – 714,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected in such a way that a pure CMIT/MIT concentration of 3,7 to less than 15 ppm is maintained. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

- 4.1.1.

Use-specific instructions for use

See general directions for use.
- 4.1.2.

Use-specific risk mitigation measures

See general directions for use.
- 4.1.3.

Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-
- 4.1.4.

Where specific to the use, the instructions for safe disposal of the product and its packaging

-
- 4.1.5.

Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-
- 4.2.

Use description

Table 2

In-can preservation of paints and coatings (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Paints and coatings
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected.The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 382,7 – 714,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 11,1 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.2.1. *Use-specific instructions for use*

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

See general directions for use.

4.2.2. *Use-specific risk mitigation measures*

See general directions for use.

4.2.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.2.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.2.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.3. **Use description**

Table 3

In-can preservation of fluids used in paper production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -

Field(s) of use	indoor use Fluids used in paper production (e.g. pigment slurries)
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 255,1 – 1 428,6 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.3.1. *Use-specific instructions for use*

See general directions for use.

4.3.2. *Use-specific risk mitigation measures*

See general directions for use.

4.3.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.3.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.3.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.4. **Use description**

Table 4

In-can preservation of fluids used in textile production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in textile production
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 127,6 – 1 428,6 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.4.1. Use-specific instructions for use

See general directions for use.

4.4.2. Use-specific risk mitigation measures

See general directions for use.

4.4.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.4.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.4.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.5. Use description

Table 5

In-can preservation of fluids used in leather production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in leather production
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 255,1 – 1 428,6 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.5.1. Use-specific instructions for use

See general directions for use.

4.5.2. Use-specific risk mitigation measures

See general directions for use.

4.5.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.5.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.5.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.6. Use description

Table 6

In-can preservation of glues and adhesives (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Glues and adhesives
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 127,6 – 1 428,6 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 30 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.6.1. *Use-specific instructions for use*

See general directions for use.

4.6.2. *Use-specific risk mitigation measures*

See general directions for use.

4.6.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.6.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.6.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.7. Use description

Table 7

In-can preservation of concrete additives and building material such as fillers/sealants, plasters and wax emulsions (PT 6) for indoor use

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Concrete additives and building material such as fillers/sealants, plasters and wax emulsions intended for indoor use.
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected during the production. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 255,1 - 714,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.7.1. *Use-specific instructions for use*

See general directions for use.

4.7.2. *Use-specific risk mitigation measures*

See general directions for use.

4.7.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.7.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.7.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.8. **Use description**

Table 8

In-can preservation of polymer dispersions/emulsions intended for use in paints and coatings, glues and adhesives, building materials and fluids used in textile production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Polymer dispersion/emulsions intended for use in paints and coatings, glues and adhesives, building materials and fluids used in textile production. Aim of application is the preservation of polymer mixtures which are consecutively used for formulation of paints, glues and further matrices (covered by PT 6).
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 127,6 - 714,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.8.1. *Use-specific instructions for use*

See general directions for use.

4.8.2. *Use-specific risk mitigation measures*

See general directions for use.

4.8.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.8.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.8.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.9. **Use description**

Table 9

In-can preservation of mineral slurries (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: -
Field(s) of use	indoor use Mineral slurries, e.g. CaCO ₃ -slurries as used for instance in paper industry and other industrial branches.
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 51,0 - 714,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 1,48 to 15 ppm. Number and timing of application: Frequency: Single application

Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.9.1. *Use-specific instructions for use*

See general directions for use.

4.9.2. *Use-specific risk mitigation measures*

See general directions for use.

4.9.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.9.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.9.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.10. **Use description**

Table 10

Preservation of cooling fluids in open recirculating systems (PT11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -

Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in open recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 25,3 – 714,3 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 151,7 - 714,3 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof shall be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast 24 hours, against algae 48 hours.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.10.1. Use-specific instructions for use

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

See general directions for use.

4.10.2. Use-specific risk mitigation measures

The use is restricted to small cooling systems with a maximum blowdown of 2 m³/h. Wastewater shall be discharged to the municipal sewer or purified in an on-site industrial sewage treatment plant including a biological treatment step. The product can only be applied when the cooling towers are equipped with drift eliminators that reduce drift with at least 99 %.

See general directions for use.

4.10.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.10.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.10.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.11. Use description

Table 11

Preservation of cooling fluids in closed systems (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in closed recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 25,3 – 714,3 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 151,7 - 714,3 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm.

	Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof shall be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast 24 hours, against algae 48 hours.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.11.1. *Use-specific instructions for use*

See general directions for use.

4.11.2. *Use-specific risk mitigation measures*

See general directions for use.

4.11.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.11.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.11.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.12. **Use description**

Table 12

Preservation of other liquids in e.g. air conditioning systems, air washers and pasteurizers (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). To protect water and other liquids in air conditioning systems, air washers, humidifiers, evaporate condensers, stationary sprinklers, water spray extinguishing systems and pasteurizers, from microbial growth.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 25,3 – 714,3 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into liquids used in air conditioning systems, air washers, humidifiers, evaporate condensers, membrane systems, stationary sprinkler, water spray extinguishing systems and pasteurizers to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm for preventive treatment against bacteria (incl. legionella), fungi and yeast, for preventive treatment against algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 151,7 - 714,3 mg product/kg matrix to be preserved. For curative treatment a pure CMIT/MIT concentration of 4,4 to 15 ppm shall be used. Number and timing of application: Frequency: Single or multiple application (number of additions depends on the state of the unit being treated) Contact time for curative use against bacteria (incl. legionella), fungi and yeast 24 hours, against algae 48 hours.
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.1.2.1. Use-specific instructions for use

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

See general directions for use.

4.12.2. *Use-specific risk mitigation measures*

See general directions for use.

4.12.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.12.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.12.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.13. **Use description**

Table 13

Preservation of wood preservative treatment solutions (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Preservation of wood preservative treatment solutions
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the wood preservative solution to be protected. Solutions of the biocidal product are automatically dosed in the wood preservative solution during the flow.
Application rate(s) and frequency	Application Rate: 1583 - 2381 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into wood preservative treatment solution to be protected leading to a pure CMIT/MIT concentration of 45,9 to 50 ppm. Number and timing of application: Frequency: Number of additions per week depends on the state of the unit being treated.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.13.1. *Use-specific instructions for use*

See general directions for use.

4.13.2. Use-specific risk mitigation measures

The product shall not be applied to wood preservative treatment solutions which are used to preserve wood intended for direct contact with water or soils (use class 4).

Freshly treated timber must be stored after treatment under shelter or on impermeable hard standing, or both, to prevent direct losses to soil, sewer or water and that any losses shall be collected for reuse or disposal.

See general directions for use.

4.13.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.13.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.13.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.14. Use description

Table 14

Preservation of membrane module solutions (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: -
Field(s) of use	indoor use Membrane module solutions. Conservation / Preservation of the water in membrane units used in water pre-treatment preparation during production shutdown after cleaning. The biocides are applied for preservation of process fluids used for non-food membrane units/systems (e.g. reverse osmosis and ultrafiltration membranes), which are widely used in the water pre-treatment preparation (non-food, non potable water, non-medical). Membrane units are used in different industries (wastewater, surface technologies, sea water desalination, etc.). These systems are recirculating.

Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: 355,2 – 809,5 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 10,3 to 17 ppm. Number and timing of application: Frequency: Single or multiple application (number of additions depends on the state of the unit being treated)
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.14.1. *Use-specific instructions for use*

During the shutdown of production, it is recommended to counteract microbiological growth inside the membrane elements. Before the shutdown, the plant shall be cleaned first, to remove all sorts of deposits like carbonates, salts, silica, organic matter or biomass. For this purpose, specialized membrane cleaners are used. After cleaning, the plant is flushed with permeate quality water until neutral pH is achieved. Only after these cleaning steps, CMIT/MIT containing water is reaching the plant and is regularly circulated by slow pumping. The biocidal product shall be added to the fill-water (permeate quality), to avoid microbiological growth during longer shut downs. During longer shut downs, the fill-water shall be regularly circulated by slow pumping and sampled to check for a microbiological re-contamination. A shift in pH can be a first indication of a re-contamination by microbiology. In this case the fill-up solution must be replaced.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

See general directions for use.

4.14.2. *Use-specific risk mitigation measures*

See general directions for use.

4.14.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.14.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.14.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.15. Use description

Table 15

Preservation of white water circuits (short circulation) in paper industry (PT 12)

Product type	PT12: Slimicides
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Preservation of white water circuits (short circulation) in paper industry. a. To protect white water circuits (short circulation) in paper machines (preventive) against bacteria. b. For curative treatment in of white water circuits (short circulation) in paper machines already affected against bacteria and fungi. Aim of the application is to prevent or combat slime formation on pipework, and surfaces of materials and equipment in the paper area.
Application method(s)	Method: dosing directly into the system (Shock / Continuous dosing) Detailed description: The biocidal product is dosed directly into the system to be protected. (Shock / Continuous dosing). Solutions of the biocidal product are automatically added by means of a dosing pump and pipes directly in the circuit (head box, mixing chest, collecting sump, broke, etc.), in intermittent or shock dosage.
Application rate(s) and frequency	Application Rate: a. Preventive: 25,5 – 714,3 mg product/kg matrix to be preserved (strongly dependent on actual system being treated). For preventive treatment (maintenance) in white water circuits (short circulation), the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,74 to 15 ppm. This refers to max. 10,6 g a.i./ton paper. b. Curative: 253,4 – 714,3 mg product/kg matrix to be preserved (strongly dependent on actual system being treated) For curative treatment (shock dose) in white water circuits (short circulation), the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 7,35 to 15 ppm. Number and timing of application: Frequency: The number of additions per week depends on the state of the unit being treated. The addition frequency of biocidal product in shock dosage is 1 to 6 times per day. Contact time for curative use: 24 hours.

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.15.1. *Use-specific instructions for use*

See general directions for use.

4.15.2. *Use-specific risk mitigation measures*

Application is only allowed in paper factories that comply with the Industrial Emission Directive 2010/75/EU where wastewater is purified in an on-site industrial sewage treatment plant including a biological treatment step in accordance to the Best Available Techniques (BAT) as prescribed in the BAT-reference document (BREF) for the production of pulp, paper and board. The effluent must be diluted at least 200 times. Paper factories that are exempted from the Industrial Emission Directive must discharge to the municipal sewer.

See general directions for use.

4.15.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.15.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.15.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.16. **Use description**

Table 16

Preservation of liquids used to treat or cut metal and glass (PT 13)

Product type	PT13: Working or cutting fluid preservatives
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use For use as a protective agent in liquids used to treat or cut metal and glass. Aim of application is to prevent microbial deterioration caused by bacteria, fungi or yeasts.

Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is added to the emulsion already formed. The biocide is added mechanical or fully automated with a timer. For tank-side addition, the biocidal products are added with an automated dosing pump and specific pipes into the collecting sump where the metal working fluids flow. The product introduction shall be done below the water level to enable a fast mixing and reduce the biocide exposure risk. The dosage is done for 30-60 minutes.
Application rate(s) and frequency	Application Rate: 817,2 - 1 428,6 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 23,7 to 30 ppm. Number and timing of application: Frequency: 1-7 times a week
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

- 4.16.1.

Use-specific instructions for use

See general directions for use.
- 4.16.2.

Use-specific risk mitigation measures

See general directions for use.
- 4.16.3.

Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-
- 4.16.4.

Where specific to the use, the instructions for safe disposal of the product and its packaging

-
- 4.16.5.

Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

5. GENERAL DIRECTIONS FOR USE OF THE META SPC 4

5.1. Instructions for use

General instructions

Preliminary tests to determine suitability, compatibility and optimal concentration are recommended prior to the use of the product in a new application. For more technical information, see the currently valid product information sheet which will be provided to customers upon request.

The product labels give instructions on the dosage of the product.

General use information**PT 6**

For the in-can preservation of washing and cleaning fluids, paints and coatings, fluids used in paper, textile and leather production, as well as glues and adhesives, concrete additives and building materials (e.g. fillers/sealants, plasters, wax emulsions), polymer dispersions/ emulsions (intended for use in paints, coatings, glues, adhesives, building materials and fluids used in textile production) and the treatment of mineral slurries.

The product shall be homogeneously incorporated into the matrix to be protected.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

PT 11

For use as a biocide for the preservation of cooling fluids in circulating systems and other liquids in e.g. air conditioning systems, air washers and pasteurizers from infestation with harmful organisms such as bacteria (including *Legionella*), fungi, yeasts and algae. Furthermore, the applications covered by PT 11 comprise the preservation of membrane module solutions and wood preservative treatment solutions.

Heavily contaminated units shall be cleaned prior to treatment.

The biocidal product is added to the water circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The microbiocidal effect begins immediately after metering.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used. Due to variation in the different systems and matrices (organic load, microbial contamination, slime forming, temperature, pH, etc.) accurate measurements by chemical and microbiological tests shall be carried out to determine the effective dose for the specific location or system.

PT 12

For use as a biocide to for the preservation of white water circuits (short circulation) in paper machines. Aim of the application is to prevent or combat slime formation (as caused by bacteria and fungi) on pipework and surfaces of materials and equipment in the paper area.

The microbiocidal effect begins immediately after metering.

The biocidal product is added to the white water circuit (short circulation) in paper machines to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

PT 13

For use as a protective agent for the preservation of liquids used to treat or cut metal and glass. Aim of application is to prevent microbial deterioration caused by bacteria, fungi or yeasts.

A ready-to-use biocidal product, without former dilution or formulation), is directly applied into the ready to use metalworking fluid cycle to control microbial deterioration and to maintain the correct function of the metalworking fluid.

The microbiocidal effect begins immediately after metering.

The biocidal product is added to the metalworking circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

5.2. Risk mitigation measures

During product handling: Wear protective gloves consistent with European Standard EN ISO 374 or equivalent, coverall (Chemical protection clothes type 6) consistent with European Standard EN ISO 13034 or equivalent and eye or face protection (recommended splash goggles or face shield) consistent with European Standard EN ISO 16321 or equivalent.

In case of C(M)IT/MIT concentrations ≥ 15 ppm: During product handling, application and post-application: Wear protective chemical resistant gloves consistent with European Standard EN ISO 374 or equivalent and a protective coverall (at least type 6) consistent with European Standard EN ISO 13034 or equivalent which is impermeable for the biocidal product.

Equipment should preferably be cleaned with water. Dilute rinsing water sufficiently before disposal via a wastewater treatment plant.

Only vent waste air to the atmosphere via suitable separators or scrubbers. No special fire or explosion protection measures necessary.

Required Personal Protection Equipment is described in Safety Data Sheet.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

First aid measures

IF SWALLOWED: Immediately rinse mouth. Give something to drink, if exposed person is able to swallow. Do NOT induce vomiting. Call 112/ambulance for medical assistance.

IF INHALED: Move to fresh air and keep at rest in a position comfortable for breathing.

If symptoms: Call 112/ambulance for medical assistance.

If no symptoms: Call a POISON CENTRE or a doctor.

IF ON SKIN: Immediately wash skin with plenty of water. Thereafter take off all contaminated clothing and wash it before reuse. Continue to wash the skin with water for 15 minutes. Call a POISON CENTRE or a doctor.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing for at least 15 minutes. Call 112/ambulance for medical assistance.

Particular environmental risks

Product containers must be carefully handled, stored and transported to prevent damage to the containers and leakage of the product into soil, air and water. The biocidal product is harmful to aquatic organisms. The product is biodegradable following dilution.

5.4. Instructions for safe disposal of the product and its packaging

Disposal information

Check whether re-use of the product in the packaging is possible. Pack or seal product waste and contaminated empty containers, label, and discharge in accordance with refuse disposal acts.

In the case of larger amounts of product, discuss with the supplier of the product in the packaging. The recipient of contaminated empty containers shall be informed of the possible risks due to product residues. In the case of disposal within the EU, use the respective valid waste code in accordance with the European Waste Catalogue (EWC). The producer of the waste is responsible for assigning sector- and process-specific waste codes to his waste in accordance with the European Waste Catalogue.

5.5. **Conditions of storage and shelf-life of the product under normal conditions of storage**

Shelf-life: 24 months.

Store in tightly sealed original containers. Protect from frost.

6. **OTHER INFORMATION**

With respect to the “Category (ies) of users” note: “Professionals (including industrial users) means trained professionals if this is required by national legislation.”

Resistance management for intended applications (PT 11, 12, 13):

- avoidance of under-dosing;
- frequent detection of efficacy and of biocide content in the industrial systems to assure maintenance of correct CMIT/MIT concentration;
- for new applications, preliminary tests to determine the suitability, compatibility and optimal application concentration are strongly recommended;
- under difficult conditions alternation of active ingredients, i.e. rotation with other biocides, also combination with other products may be useful.

Full titles of EN standards and legislation referred to in section 5.2:

EN ISO 374 – Protective gloves against dangerous chemicals and micro-organisms.

EN ISO 13034 - Protective clothing against liquid chemicals - Performance requirements for chemical protective clothing offering limited protective performance against liquid chemicals (Type 6 and Type PB [6] equipment)

EN ISO 16321 – Eye and face protection for occupational use

7. **THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 4**

7.1. **Trade name(s), authorisation number and specific composition of each individual product**

Trade name(s)			Preventol D7 Plus	Market area: EU		
Authorisation number			EU-0031652-0016 1-4			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		3,6 % (w/w)	

7.2. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Ferrocid 8583	Market area: EU		
Authorisation number			EU-0031652-0017 1-4			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		3,6 % (w/w)	

7.3. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 04-03	Market area: EU		
Authorisation number			EU-0031652-0018 1-4			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		3,6 % (w/w)	

7.4. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 04-04	Market area: EU		
Authorisation number			EU-0031652-0019 1-4			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		4,1 % (w/w)	

1. META SPC 5 ADMINISTRATIVE INFORMATION

1.1. Meta SPC 5 identifier

Identifier	Meta SPC: Meta SPC 5
------------	----------------------

1.2. Suffix to the authorisation number

Number	1-5
--------	-----

1.3. Product type(s)

Product type(s)	PT11: Preservatives for liquid-cooling and processing systems
-----------------	---

2. META SPC 5 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 5

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		1 - 1,7 % (w/w)

2.2. Type(s) of formulation of the meta SPC 5

Formulation type(s)	AL Any other liquid
---------------------	---------------------

3. HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 5

Hazard statements	H314: Causes severe skin burns and eye damage. H317: May cause an allergic skin reaction. H410: Very toxic to aquatic life with long lasting effects. EUH071: Corrosive to the respiratory tract.
Precautionary statements	P272: Contaminated work clothing should not be allowed out of the workplace. P273: Avoid release to the environment. P280: Wear protective gloves. P280: Wear protective clothing. P280: Wear eye protection. P280: Wear face protection. P301+P330+P331: IF SWALLOWED: rinse mouth. Do NOT induce vomiting.

	P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water [or shower]. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P362+P364: Take off contaminated clothing and wash it before reuse. P405: Store locked up. P501: Dispose of contents in accordance with all local, regional, national and international regulations.. P501: Dispose of container in accordance with all local, regional, national and international regulations.. P391: Collect spillage.
--	---

4. AUTHORISED USE(S) OF THE META SPC

4.1. Use description

Table 1

Preservation of cooling fluids in open recirculation systems (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in open recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.

Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 73,5 – 2500 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 440,0 - 2500 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. Contact time for curative use against bacteria (incl. legionella), fungi and yeast 24 hours, against algae 48 hours.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.1.1. Use-specific instructions for use

See general directions for use.

4.1.2. Use-specific risk mitigation measures

The use is restricted to small cooling systems with a maximum blowdown of 2 m³/h. Waste water shall be discharged to the municipal sewer or purified in an on-site industrial sewage treatment plant including a biological treatment step. The product shall only be applied when the cooling towers are equipped with drift eliminators that reduce drift with at least 99 %.

See general directions for use.

4.1.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.1.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.1.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.2. Use description

Table 2

Preservation of cooling fluids in closed systems (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in closed recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 73,5 – 2500 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 440,0 - 2500 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. Contact time for curative use against bacteria (incl. legionella), fungi and yeast 24 hours, against algae 48 hours.

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

- 4.2.1.

Use-specific instructions for use

See general directions for use.
- 4.2.2.

Use-specific risk mitigation measures

See general directions for use.
- 4.2.3.

Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-
- 4.2.4.

Where specific to the use, the instructions for safe disposal of the product and its packaging

-
- 4.2.5.

Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

5. GENERAL DIRECTIONS FOR USE OF THE META SPC 5

5.1. Instructions for use

General instructions

Preliminary tests to determine suitability, compatibility and optimal concentration are recommended prior to the use of the product in a new application. For more technical information, see the currently valid product information sheet which will be provided to customers upon request.

The product labels give instructions on the dosage of the product.

General use information

PT11

For use as a biocide for the preservation of cooling fluids in circulating systems from infestation with harmful organisms such as bacteria (including Legionella), fungi, yeasts and algae.

Heavily infested units shall be cleaned prior to treatment.

The biocidal product is added to the water circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The microbiocidal effect begins immediately after metering. The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

5.2. Risk mitigation measures

During product handling: Wear protective gloves consistent with European Standard EN ISO 374 or equivalent, coverall (Chemical protection clothes type 6) consistent with European Standard EN ISO 13034 or equivalent and eye or face protection (recommended splash goggles or face shield) consistent with European Standard EN ISO 16321 or equivalent.

In case of C(M)IT/MIT concentrations ≥ 15 ppm: During product handling, application and post-application: Wear protective chemical resistant gloves consistent with European Standard EN ISO 374 or equivalent and a protective coverall (at least type 6) consistent with European Standard EN ISO 13034 or equivalent which is impermeable for the biocidal product.

Equipment should preferably be cleaned with water. Dilute rinsing water sufficiently before disposal via a wastewater treatment plant.

Only vent waste air to the atmosphere via suitable separators or scrubbers. No special fire or explosion protection measures necessary.

Required Personal Protection Equipment is described in Safety Data Sheet.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

First aid measures

IF SWALLOWED: Immediately rinse mouth. Give something to drink, if exposed person is able to swallow. Do NOT induce vomiting. Call 112/ambulance for medical assistance.

IF INHALED: Move to fresh air and keep at rest in a position comfortable for breathing.

If symptoms: Call 112/ambulance for medical assistance.

If no symptoms: Call a POISON CENTRE or a doctor.

IF ON SKIN: Immediately wash skin with plenty of water. Thereafter take off all contaminated clothing and wash it before reuse. Continue to wash the skin with water for 15 minutes. Call a POISON CENTRE or a doctor.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing for at least 15 minutes. Call 112/ambulance for medical assistance.

Particular environmental risks

Product containers must be carefully handled, stored and transported to prevent damage to the containers and leakage of the product into soil, air and water. The biocidal product is harmful to aquatic organisms. The product is biodegradable following dilution.

5.4. Instructions for safe disposal of the product and its packaging

Disposal information

Check whether re-use of the product in the packaging is possible. Pack or seal product waste and contaminated empty containers, label, and discharge in accordance with refuse disposal acts.

In the case of larger amounts of product, discuss with the supplier of the product in the packaging. The recipient of contaminated empty containers must be informed of the possible risks due to product residues. In the case of disposal within the EU, use the respective valid waste code in accordance with the European Waste Catalogue (EWC). The producer of the waste is responsible for assigning sector- and process-specific waste codes to his waste in accordance with the European Waste Catalogue.

5.5. **Conditions of storage and shelf-life of the product under normal conditions of storage**

Shelf-life: 24 months.

Store in tightly sealed original containers. Protect from frost.

6. **OTHER INFORMATION**

With respect to the “Category (ies) of users” note: “Professionals (including industrial users) means trained professionals if this is required by national legislation.”

Resistance management for intended applications (PT 11):

- avoidance of under-dosing;
- frequent detection of efficacy and of biocide content in the industrial systems to assure maintenance of correct CMIT/MIT concentration;
- for new applications, preliminary tests to determine the suitability, compatibility and optimal application concentration are strongly recommended;
- under difficult conditions alternation of active ingredients, i.e. rotation with other biocides, also combination with other products may be useful.

Full titles of EN standards and legislation referred to in section 5.2:

EN ISO 374 – Protective gloves against dangerous chemicals and micro-organisms.

EN ISO 13034 - Protective clothing against liquid chemicals - Performance requirements for chemical protective clothing offering limited protective performance against liquid chemicals (Type 6 and Type PB [6] equipment)

7. **THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 5**

7.1. **Trade name(s), authorisation number and specific composition of each individual product**

Trade name(s)			Preservative 05-01	Market area: EU	
Authorisation number			EU-0031652-0020 1-5		
Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		1,3 % (w/w)

1. META SPC 6 ADMINISTRATIVE INFORMATION

1.1. Meta SPC 6 identifier

Identifier	Meta SPC: Meta SPC 6
------------	----------------------

1.2. Suffix to the authorisation number

Number	1-6
--------	-----

1.3. Product type(s)

Product type(s)	PT06: Preservatives for products during storage PT11: Preservatives for liquid-cooling and processing systems PT12: Slimicides PT13: Working or cutting fluid preservatives
-----------------	--

2. META SPC 6 COMPOSITION

2.1. Qualitative and quantitative information on the composition of the meta SPC 6

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro-2-methyl-2h-isothiazol-3-one and 2-methyl-2h-isothia zol-3-one (3:1)	Active substance	55965-84-9		1,7 - 2,5 % (w/w)

2.2. Type(s) of formulation of the meta SPC 6

Formulation type(s)	AL Any other liquid
---------------------	---------------------

3. HAZARD AND PRECAUTIONARY STATEMENTS OF THE META SPC 6

Hazard statements	H314: Causes severe skin burns and eye damage. H317: May cause an allergic skin reaction. H410: Very toxic to aquatic life with long lasting effects. EUH071: Corrosive to the respiratory tract.
-------------------	--

Precautionary statements	P272: Contaminated work clothing should not be allowed out of the workplace. P273: Avoid release to the environment. P280: Wear protective gloves. P280: Wear protective clothing. P280: Wear eye protection. P280: Wear face protection. P301+P330+P331: IF SWALLOWED: rinse mouth. Do NOT induce vomiting. P310: Immediately call a doctor. P310: Immediately call a POISON CENTER. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water [or shower]. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P310: Immediately call a POISON CENTER. P310: Immediately call a doctor. P362+P364: Take off contaminated clothing and wash it before reuse. P405: Store locked up. P501: Dispose of contents in accordance with all local, regional, national and international regulations. P501: Dispose of container in accordance with all local, regional, national and international regulations.. P391: Collect spillage.
--------------------------	---

4. **AUTHORISED USE(S) OF THE META SPC**

4.1. **Use description**

Table 1

In-can preservation of washing and cleaning fluids (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Washing and cleaning fluids

Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected.The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 261 – 1 500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected in such a way that a pure CMIT/MIT concentration of 3,7 to less than 15 ppm is maintained. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

- 4.1.1.

Use-specific instructions for use

See general directions for use.
- 4.1.2.

Use-specific risk mitigation measures

See general directions for use.
- 4.1.3.

Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-
- 4.1.4.

Where specific to the use, the instructions for safe disposal of the product and its packaging

-
- 4.1.5.

Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-
- 4.2.

Use description

Table 2

In-can preservation of paints and coatings (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Paints and coatings
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 782 – 1500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 11,1 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.2.1. Use-specific instructions for use

See general directions for use.

4.2.2. Use-specific risk mitigation measures

See general directions for use.

4.2.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.2.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.2.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.3. Use description

Table 3

In-can preservation of fluids used in paper production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in paper production (e.g. pigment slurries)
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 521 – 3000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.3.1. Use-specific instructions for use

See general directions for use.

4.3.2. Use-specific risk mitigation measures

See general directions for use.

4.3.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.3.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.3.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.4. Use description

Table 4

In-can preservation of fluids used in textile production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in textile production
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 261 – 3000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.4.1. Use-specific instructions for use

See general directions for use.

4.4.2. Use-specific risk mitigation measures

See general directions for use.

4.4.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.4.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.4.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.5. Use description

Table 5

In-can preservation of fluids used in leather production (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Fluids used in leather production
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 521 – 3000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 30 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.5.1. *Use-specific instructions for use*

See general directions for use.

4.5.2. *Use-specific risk mitigation measures*

See general directions for use.

4.5.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.5.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.5.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.6. Use description

Table 6

In-can preservation of glues and adhesives (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Glues and adhesives
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 261 – 3000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 30 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.6.1. *Use-specific instructions for use*

See general directions for use.

4.6.2. *Use-specific risk mitigation measures*

See general directions for use.

4.6.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.6.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.6.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.7. **Use description**

Table 7

In-can preservation of concrete additives and building material such as fillers/sealants, plasters and wax emulsions (PT 6) for indoor use

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Concrete additives and building material such as fillers/sealants, plasters and wax emulsions intended for indoor use
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 521 - 1500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 7,4 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.7.1. *Use-specific instructions for use*

See general directions for use.

4.7.2. *Use-specific risk mitigation measures*

See general directions for use.

4.7.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.7.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.7.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.8. **Use description**

Table 8

In-can preservation of polymer dispersions/emulsions intended for use in paints and coatings, glues and adhesives, building materials, fluids used in textile production and plant protection products (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Polymer dispersions/emulsions intended for use in paints and coatings, glues and adhesives, building materials, fluids used in textile production and plant protection products. Aim of application is the preservation of polymer mixtures which are consecutively used for formulation of paints, glues and further matrices (covered by PT 6).

Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected. The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 261 - 1 500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 3,7 to 15 ppm. End products which are used by the general public shall contain less than 15 ppm active ingredient. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.8.1. *Use-specific instructions for use*

See general directions for use.

4.8.2. *Use-specific risk mitigation measures*

See general directions for use.

4.8.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.8.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.8.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.9. **Use description**

Table 9

In-can preservation of mineral slurries (PT 6)

Product type	PT06: Preservatives for products during storage
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: -
Field(s) of use	indoor use Mineral slurries, e.g. CaCO3-slurries as used for instance in paper industry and other industrial branches.
Application method(s)	Method: Manual and automatic dosing Detailed description: The biocidal product shall be homogeneously incorporated into the matrix to be protected.The biocidal product is dosed in the mixing vessel during the waterborne system production, usually as a last ingredient.
Application rate(s) and frequency	Application Rate: 104 - 1500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration 1,48 to 15 ppm. Number and timing of application: Frequency: Single application
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

- 4.9.1. *Use-specific instructions for use*
See general directions for use.
- 4.9.2. *Use-specific risk mitigation measures*
See general directions for use.
- 4.9.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*
-
- 4.9.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*
-
- 4.9.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*
-
- 4.10. **Use description**

Table 10

Preservation of cooling fluids in open recirculating systems (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-

Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in open recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 51,8 – 1 500 mg product/kg matrix to be preserved. For preventive treatment in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm. b. Curative: 309,9 - 1 500 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof should be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast: 24 hours, against algae: 48.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.10.1. Use-specific instructions for use

See general directions for use.

4.10.2. Use-specific risk mitigation measures

The use is restricted to small cooling systems with a maximum blowdown of 2 m³/h. Waste water must be discharged to the municipal sewer or purified in an on-site industrial sewage treatment plant including a biological treatment step. The product can only be applied when the cooling towers are equipped with drift eliminators that reduce drift with at least 99 %.

See general directions for use.

4.10.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.10.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.10.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.11. Use description

Table 11

Preservation of cooling fluids in closed systems (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). a. Fluids in closed recirculating liquid-cooling systems with discharge to the municipal sewer (preventive). b. For curative treatment in cooling water circuits already affected.

Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 51,8 – 1500 mg product/kg matrix to be preserved. For preventive treatment against bacteria (incl. legionella), fungi and yeast in cooling water circuits, the biocidal product shall be incorporated into product to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm, for algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 309,9 - 1500 mg product/kg matrix to be preserved. For curative treatment in cooling water circuits, the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 4,4 to 15 ppm. Number and timing of application: Frequency: a. Preventive: Number of additions per week depends on the state of the unit being treated. b. Curative: 2 - 3 times a week until an acceptable microbial contamination is reached. An acceptable microbial contamination and the measurement thereof should be in line with the users 'hygiene management system' in place. Contact time for curative use against bacteria (incl. legionella), fungi and yeast: 24 hours, against algae: 48 hours.
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.11.1. *Use-specific instructions for use*

See general directions for use.

4.11.2. *Use-specific risk mitigation measures*

See general directions for use.

4.11.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.11.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.11.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.12. Use description

Table 12

Preservation of other liquids in e.g. air conditioning systems, air washers and pasteurizers (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: algae Development stage: - Scientific name: - Common name: Legionella Development stage: -
Field(s) of use	indoor use Dosing stations may be placed outdoor (roofed or in container). To protect water and other liquids in air conditioning systems, air washers, humidifiers, evaporate condensers, stationary sprinklers, water spray extinguishing systems and pasteurizers, from microbial growth.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: a. Preventive: 51,8 – 1500 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into liquids used in air conditioning systems, air washers, humidifiers, evaporate condensers, membrane systems, stationary sprinkler, water spray extinguishing systems and pasteurizers to be protected leading to a pure CMIT/MIT concentration of 0,735 to 15 ppm for preventive treatment against bacteria (incl. legionella), fungi and yeast, for preventive treatment against algae a pure CMIT/MIT concentration of 1,42 to 15 ppm shall be used. b. Curative: 309,9 - 1500 mg product/kg matrix to be preserved. For curative treatment a pure CMIT/MIT concentration of 4,4 to 15 ppm shall be used. Number and timing of application: Frequency: Single or multiple application (number of additions depends on the state of the unit being treated) Contact time for curative use against bacteria (incl. legionella), fungi and yeast 24 hours, against algae 48 hours.

Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.12.1. *Use-specific instructions for use*

See general directions for use.

4.12.2. *Use-specific risk mitigation measures*

See general directions for use.

4.12.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.12.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.12.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.13. **Use description**

Table 13

Preservation of membrane module solutions (PT 11)

Product type	PT11: Preservatives for liquid-cooling and processing systems
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: - Scientific name: - Common name: yeasts Development stage: -
Field(s) of use	indoor use Membrane module solutions. Conservation / Preservation of the water in membrane units used in water pre-treatment preparation during production shutdown after cleaning. The biocides are applied for preservation of process fluids used for non-food membrane units/systems (e.g. reverse osmosis and ultrafiltration membranes), which are widely used in the water pre-treatment preparation (non-food, non potable water, non-medical). Membrane units are used in different industries (wastewater, surface technologies, sea water desalination, etc.). These systems are recirculating.

Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is dosed directly into the system to be protected. Solutions of biocidal product are automatically dosed in the water of the circuit during the flow.
Application rate(s) and frequency	Application Rate: 725,4 – 1700 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected, leading to a pure CMIT/MIT concentration of 10,3 to 17 ppm. Number and timing of application: Frequency: Single or multiple application (number of additions depends on the state of the unit being treated)
Category(ies) of users	industrial
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.1.3.1. *Use-specific instructions for use*

During the shutdown of production, it is recommended to counteract microbiological growth inside the membrane elements. Before the shutdown, the plant shall be cleaned first, to remove all sorts of deposits like carbonates, salts, silica, organic matter or biomass. For this purpose, specialized membrane cleaners are used. After cleaning, the plant is flushed with permeate quality water until neutral pH is achieved. Only after these cleaning steps, CMIT/MIT containing water is reaching the plant and is regularly circulated by slow pumping. The biocidal product shall be added to the fill-water (permeate quality), to avoid microbiological growth during longer shut downs. During longer shut downs, the fill-water shall be regularly circulated by slow pumping and sampled to check for a microbiological re-contamination. A shift in pH can be a first indication of a re-contamination by microbiology. In this case the fill-up solution shall be replaced.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

See general directions for use.

4.1.3.2. *Use-specific risk mitigation measures*

See general directions for use.

4.1.3.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.1.3.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.1.3.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

4.14. Use description

Table 14

Preservation of white water circuits (short circulation) in paper industry (PT 12)

Product type	PT12: Slimicides
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: fungi Development stage: -
Field(s) of use	indoor use Preservation of white water circuits (short circulation) in paper industry. a. To protect white water circuits (short circulation) in paper machines (preventive) against bacteria. b. For curative treatment in of white water circuits (short circulation) in paper machines already affected against bacteria and fungi. Aim of the application is to prevent or combat slime formation on pipework, and surfaces of materials and equipment in the paper area.
Application method(s)	Method: Dosing directly into the system (Shock / Continuous dosing). Detailed description: The biocidal product is dosed directly into the system to be protected. (Shock / Continuous dosing). Solutions of the biocidal product are automatically added by means of a dosing pump and pipes directly in the circuit (head box, mixing chest, collecting sump, broke, etc.), in intermittent or shock dosage.
Application rate(s) and frequency	Application Rate: a. Preventive: 52,15 – 1500 mg product/kg matrix to be preserved (strongly dependent on actual system being treated). For preventive treatment (maintenance) in white water circuits (short circulation), the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 0,74 to 15 ppm. This refers to max. 10,6 g a.i./ton paper. b. Curative: 517,6 – 1500 mg product/kg matrix to be preserved (strongly dependent on actual system being treated). For curative treatment (shock dose) in white water circuits (short circulation), the biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 7,35 – 15 ppm. Number and timing of application: Frequency: The number of additions per week depends on the state of the unit being treated. The addition frequency of biocidal product in shock dosage is 1 to 6 times per day. Contact time for curative use: 24 hours.

Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.14.1. Use-specific instructions for use

See general directions for use.

4.14.2. Use-specific risk mitigation measures

Application is only allowed in paper factories that comply with the Industrial Emission Directive 2010/75/EU where wastewater is purified in an on-site industrial sewage treatment plant including a biological treatment step in accordance to the Best Available Techniques (BAT) as prescribed in the BAT-reference document (BREF) for the production of pulp, paper and board. The effluent must be diluted at least 200 times. Paper factories that are exempted from the Industrial Emission Directive must discharge to the municipal sewer.

See general directions for use.

4.14.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

-

4.14.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.14.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.15. Use description

Table 15

Preservation of liquids used to treat or cut metal and glass (PT 13)

Product type	PT13: Working or cutting fluid preservatives
Where relevant, an exact description of the authorised use	-
Target organism(s) (including development stage)	Scientific name: - Common name: bacteria Development stage: - Scientific name: - Common name: yeasts Development stage: - Scientific name: - Common name: fungi Development stage: -

Field(s) of use	indoor use For use as a protective agent in liquids used to treat or cut metal and glass. Aim of application is to prevent microbial deterioration caused by bacteria, fungi or yeasts.
Application method(s)	Method: dosing directly into the system Detailed description: The biocidal product is added to the emulsion already formed. The biocide is added mechanical or fully automated with a timer. For tank-side addition, the biocidal products are added with an automated dosing pump and specific pipes into the collecting sump where the metal working fluids flow. The product introduction shall be done below the water level to enable a fast mixing and reduce the biocide exposure risk. The dosage is done for 30-60 minutes.
Application rate(s) and frequency	Application Rate: 1 669,0 - 3 000 mg product/kg matrix to be preserved. The biocidal product shall be incorporated into the matrix to be protected leading to a pure CMIT/MIT concentration of 23,7 to 30 ppm. Number and timing of application: Frequency: 1-7 times a week
Category(ies) of users	industrial professional
Pack sizes and packaging material	— Bottle (50 – 1 500 ml), HDPE, opaque — Jerry can (5, 10, 20, 25, 30 and 60 litre), HDPE, opaque — Drum (206 and 220 litre), HDPE, opaque — Intermediate bulk container (IBC) (1 000 litre), HDPE, opaque

4.15.1. *Use-specific instructions for use*

See general directions for use.

4.15.2. *Use-specific risk mitigation measures*

See general directions for use.

4.15.3. *Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*

-

4.15.4. *Where specific to the use, the instructions for safe disposal of the product and its packaging*

-

4.15.5. *Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage*

-

5. GENERAL DIRECTIONS FOR USE OF THE META SPC 6

5.1. Instructions for use

General instructions

Preliminary tests to determine suitability, compatibility and optimal concentration are recommended prior to the use of the product in a new application. For more technical information, see the currently valid product information sheet which will be provided to customers upon request.

The product labels give instructions on the dosage of the product.

General use information

PT 6

For the in-can preservation of washing and cleaning fluids, paints and coatings, fluids used in paper, textile and leather production, as well as glues and adhesives, concrete additives and building materials (e.g. fillers/sealants, plasters, wax emulsions), polymer dispersions/ emulsions (intended for use in paints, coatings, glues, adhesives, building materials and fluids used in textile production) and the treatment of mineral slurries.

The product shall be homogeneously incorporated into the matrix to be protected.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

PT 11

For use as a biocide for the preservation of cooling fluids in circulating systems and other liquids in e.g. air conditioning systems, air washers and pasteurizers from infestation with harmful organisms such as bacteria (including Legionella), fungi, yeasts and algae. Furthermore, the applications covered by PT 11 comprise the preservation of membrane module solutions.

Heavily contaminated units shall be cleaned prior to treatment.

The biocidal product is added to the water circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The microbiocidal effect begins immediately after metering.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used. Due to variation in the different systems and matrices (organic load, microbial contamination, slime forming, temperature, pH, etc.) accurate measurements by chemical and microbiological tests shall be carried out to determine the effective dose for the specific location or system.

PT 12

For use as a biocide for the preservation of white water circuits (short circulation) in paper machines. Aim of the application is to prevent or combat slime formation (as caused by bacteria and fungi) on pipework and surfaces of materials and equipment in the paper area.

The microbiocidal effect begins immediately after metering.

The biocidal product is added to the white water circuit (short circulation) in paper machines to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

PT 13

For use as a protective agent for the preservation of liquids used to treat or cut metal and glass. Aim of application is to prevent microbial deterioration caused by bacteria, fungi or yeasts.

A ready-to-use biocidal product, without former dilution or formulation), is directly applied into the ready to use metalworking fluid cycle to control microbial deterioration and to maintain the correct function of the metalworking fluid.

The microbiocidal effect begins immediately after metering

The biocidal product is added to the metalworking circuit to be treated at a point at which a rapid and even distribution of the biocide can be ensured.

The dose strongly depends on the formulation and intended use of the product to which the preservative is added. Therefore, the user shall determine dosage requirements for their specific matrix/system to be preserved. The lowest effective dose shall be used.

5.2. Risk mitigation measures

During product handling: Wear protective gloves consistent with European Standard EN ISO 374 or equivalent, coverall (Chemical protection clothes type 6) consistent with European Standard EN ISO 13034 or equivalent and eye or face protection (recommended splash goggles or face shield) consistent with European Standard EN ISO 16321 or equivalent).

In case of C(M)IT/MIT concentrations ≥ 15 ppm: During product handling, application and post-application: Wear protective chemical resistant gloves consistent with European Standard EN ISO 374 or equivalent and a protective coverall (at least type 6) consistent with European Standard EN 13034 or equivalent which is impermeable for the biocidal product.

Equipment should preferably be cleaned with water. Dilute rinsing water sufficiently before disposal via a wastewater treatment plant.

Only vent waste air to the atmosphere via suitable separators or scrubbers. No special fire or explosion protection measures necessary.

Required Personal Protection Equipment is described in Safety Data Sheet.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment*First aid measures*

IF SWALLOWED: Immediately rinse mouth. Give something to drink, if exposed person is able to swallow. Do NOT induce vomiting. Call 112/ambulance for medical assistance.

IF INHALED: Move to fresh air and keep at rest in a position comfortable for breathing.

If symptoms: Call 112/ambulance for medical assistance.

If no symptoms: Call a POISON CENTRE or a doctor.

IF ON SKIN: Immediately wash skin with plenty of water. Thereafter take off all contaminated clothing and wash it before reuse. Continue to wash the skin with water for 15 minutes. Call a POISON CENTRE or a doctor.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing for at least 15 minutes. Call 112/ambulance for medical assistance.

Particular environmental risks

Product containers must be carefully handled, stored and transported to prevent damage to the containers and leakage of the product into soil, air and water. The biocidal product is harmful to aquatic organisms. The product is biodegradable following dilution.

5.4. **Instructions for safe disposal of the product and its packaging**

Disposal information

Check whether re-use of the product in the packaging is possible. Pack or seal product waste and contaminated empty containers, label, and discharge in accordance with refuse disposal acts.

In the case of larger amounts of product in the packaging, discuss with the supplier of the product. The recipient of contaminated empty containers must be informed of the possible risks due to product residues. In the case of disposal within the EU, use the respective valid waste code in accordance with the European Waste Catalogue (EWC). The producer of the waste is responsible for assigning sector- and process-specific waste codes to his waste in accordance with the European Waste Catalogue.

5.5. **Conditions of storage and shelf-life of the product under normal conditions of storage**

Shelf-life: 24 months.

Store in tightly sealed original containers.

Protect from frost

6. **OTHER INFORMATION**

With respect to the “Category (ies) of users” note: “Professionals (including industrial users) means trained professionals if this is required by national legislation.”

Resistance management for intended applications (PT 11, 12, 13):

- avoidance of under-dosing;
- frequent detection of efficacy and of biocide content in the industrial systems to assure maintenance of correct CMIT/MIT concentration;
- for new applications, preliminary tests to determine the suitability, compatibility and optimal application concentration are strongly recommended;
- under difficult conditions alternation of active ingredients, i.e. rotation with other biocides, also combination with other products may be useful.

Full titles of EN standards and legislation referred to in section 5.2:

EN ISO 374 – Protective gloves against dangerous chemicals and micro-organisms.

EN ISO 13034 - Protective clothing against liquid chemicals - Performance requirements for chemical protective clothing offering limited protective performance against liquid chemicals (Type 6 and Type PB [6] equipment)

7. **THIRD INFORMATION LEVEL: INDIVIDUAL PRODUCTS IN THE META SPC 6**

7.1. **Trade name(s), authorisation number and specific composition of each individual product**

Trade name(s)	Preventol D7 CF	Market area: EU
	Kathon® CMI 1.5 CF	Market area: EU
Authorisation number	EU-0031652-0021 1-6	

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,4 % (w/w)

7.2. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preventol D7 CF2	Market area: EU		
Authorisation number			EU-0031652-0022 1-6			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,5 % (w/w)	

7.3. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Kurita F-5106	Market area: EU		
Authorisation number			EU-0031652-0023 1-6			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)	
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,4 % (w/w)	

7.4. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)	Aqualead BC 08 LA	Market area: EU
Authorisation number	EU-0031652-0024 1-6	

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,5 % (w/w)

7.5. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)			Preservative 06-05	Market area: EU	
Authorisation number			EU-0031652-0025 1-6		
Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,4 % (w/w)

7.6. Trade name(s), authorisation number and specific composition of each individual product

Trade name(s)		Preservative 06-06	Market area: EU		
Authorisation number		EU-0031652-0026 1-6			
Common name	IUPAC name	Function	CAS number	EC number	Content (%)
C(M)IT/MIT (3:1)	Reaction mass of 5- chloro- 2-methyl-2h- isothiazol- 3-one and 2-methyl- 2h-isothia zol- 3-one (3:1)	Active substance	55965-84-9		2,5 % (w/w)